

## Adeno-Associated Virus-Based Gene Therapy for CNS Diseases

Michaël Hocquemiller,<sup>1,\*</sup> Laura Giersch,<sup>1,†</sup> Mickael Audrain,<sup>2–4,†</sup> Samantha Parker,<sup>1</sup> and Nathalie Cartier<sup>3,4,\*</sup>

<sup>1</sup>Lysogene, Neuilly sur Seine, France; <sup>2</sup>Université Paris Descartes, Paris, France; <sup>3</sup>INSERM UMR1169, Université Paris-Sud, Université Paris-Saclay, Orsay, France; and <sup>4</sup>CEA, DSV, I<sup>2</sup>BM, MIRCen, Fontenay-aux-Roses, France.

<sup>†</sup>L.G. and M.A. contributed equally to this work.

Gene therapy is at the cusp of a revolution for treating a large spectrum of CNS disorders by providing a durable therapeutic protein via a single administration. Adeno-associated virus (AAV)-mediated gene transfer is of particular interest as a therapeutic tool because of its safety profile and efficiency in transducing a wide range of cell types. The purpose of this review is to describe the most notable advancements in preclinical and clinical research on AAV-based CNS gene therapy and to discuss prospects for future development based on a new generation of vectors and delivery.

### INTRODUCTION

NEUROLOGICAL DISORDERS are among the most difficult pathologies to treat because of the limited access to brain structures, which are protected by physical barriers such as the blood–brain barrier (BBB); the complexity of the CNS; and cell interactions. Gene therapy is an alternative to traditional pharmacological approaches that has made important advances over the last decade in treating the nervous system.

Experimental use of gene therapy for brain tumors has increasing promise, with a multitude of research programs using recombinant adeno-associated viral (rAAV) gene therapy with systemic or direct administration of therapeutic agents into the tumor site. This work is not considered here, as the goal to influence the proliferation of abnormal cells is not within the scope of this review. The aim of this review is to provide a thorough summary of the most notable advancements in preclinical and clinical research on rAAV gene delivery in treating neurodegenerative, genetic, and acquired diseases affecting the nervous system.

AAV is a small, nonenveloped virus that has been the subject of intense research interest from the gene therapy field. rAAVs have demonstrated remarkable evidence of efficacy and safety

in a large number of animal models. They have become the most commonly used gene therapy vectors for the CNS<sup>1</sup> because of their safety, nonpathogenic nature, and ability to infect dividing and quiescent cells *in vivo*, particularly neurons. rAAVs have also demonstrated long-term gene expression *in vivo*.<sup>2</sup>

Initial preclinical proofs of concept used first-generation vectors based on AAV serotype 2. They moved from the conceptual stage to clinical trials for several inherited and acquired diseases, such as Parkinson's, Batten's, and Canavan's disease<sup>3</sup> (Table 1). Since then, several other AAV serotypes have been isolated and engineered. AAV strains with improved tissue tropisms and biodistribution have been extensively characterized in animal models. The recombinant genome of a given serotype can be packaged into the capsid of another serotype (i.e., rAAV2/5 contains the AAV2 recombinant genome packaged in the capsid proteins encoded by the *cap* gene of AAV5<sup>4</sup>). The AAV serotypes most studied in the CNS have been serotypes 1, 2, 5, 8, 9, and recombinant human (rh)10.<sup>5–8</sup> The effectiveness of a serotype depends on the brain region, the species, and the targeted cell type. These serotypes efficiently transduce neurons;

\*Correspondence: Dr. Nathalie Cartier, INSERM UMR1169, Université Paris-Sud, Université Paris-Saclay, Orsay 94100, France. E-mail: nathalie.cartier@inserm.fr; or Dr. Michaël Hocquemiller, Lysogene, 18–20 rue Jacques Dulud, Neuilly sur Seine 92200, France. E-mail: michael.hocquemiller@lysogene.com

© Michaël Hocquemiller, et al., 2016; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons Attribution Noncommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

**Table 1. Clinical Trials**

	Injection site	Disease	Clinical trial	Inclusion	Serotype	Transgene	Promoter <sup>a</sup>	Dose, min vg	Dose, max vg	Volume, $\mu$ L	Speed, $\mu$ L/min	IS	Status	Identifier	Ref.	
Intracerebral	WM (n=6)	Can	Phase I	13	2	ASP	NSE	$9 \times 10^{11}$		900	2	NA	C	NA	15	
	WM (n=12)	LINCF	Phase I	11	2	CLN2	CAG	$1.8 \times 10^{12}$ – $3.2 \times 10^{12}$		600	2	NA	C	NCT00151216	17	
	WM (n=12)	LINCF	Phase I/II	16	rh10	CLN2	CAG	$2.85 \times 10^{11}$ – $9 \times 10^{11}$		1800	2	NA	O	NCT01414985	NA	
	WM (n=12)	MPS IIIA	Phase I/II	4	rh10	SGSH	PGK	$7.2 \times 10^{11}$		720	0.5	Y	C	NCT01474343	16	
	WM (n=12)/ Cer (n=4)	MPS IIIB	Phase I/II	4	5	NAGLU	PGK	$4 \times 10^{12}$		960	0.5	Y	O	ISRCTN19853672	NA	
	WM (n=12)	MLD	Phase I/II	5	rh10	ARSA	CAG	$1 \times 10^{12}$ – $4 \times 10^{12}$		NA	NA	NA	O	NCT01801709	NA	
	StN (n=2)	Par	Phase II	16	2	GAD	CAG	$2 \times 10^{12}$		70	0.23	NA	C	NCT00643890	162	
	Str (n=4)	Par	Phase I	10	2	AADC	CMV	$9 \times 10^{10}$ – $3 \times 10^{11}$		200	1	N	C	NCT00229736	163	
	Put (n=8)	Par	Phase I&II	70	2	NTN (CERE-120)	CAG	$1.3 \times 10^{11}$ – $5.4 \times 10^{11}$		80	2	NA	C	NCT00252850 NCT00400634	106,164	
	Put (n=6)/ SN (n=4)	Par	Phase I/II	57	2	NTN (CERE-120)	CAG	$9.4 \times 10^{11}$ – $2.4 \times 10^{12}$		360	2/3	NA	O	NCT00985517	165	
	Str (n=2)	Par	Phase I	24	2	GDNF	CMV	$9 \times 10^{10}$ – $3 \times 10^{12}$		NA	NA	NA	O	NCT01621581	NA	
	Str (n=2)	Par	Phase I	10	2	AADC	NA	$7.5 \times 10^{11}$ – $1.5 \times 10^{12}$		NA	NA	NA	O	NCT01973543	NA	
	Put (n=4)	Par	Phase I/II	6	NA	AADC	NA	$3 \times 10^{11}$ – $9 \times 10^{11}$		200/600	3	NA	O	NCT02418598	NA	
	Put (n=2)	Par	Phase I	10	2	AADC	NA	NA		NA	NA	NA	O	NCT01395641	NA	
	NBM (n=4/6)	Alz	Phase I	10	2	NGF (CERE-110)	CAG	$1.2 \times 10^{10}$ – $1.2 \times 10^{11}$		40/80	2	NA	C	NCT00087789	79	
	NA	Alz	Phase II	25	2	NGF (CERE-110)	CAG	$2 \times 10^{11}$		NA	NA	NA	NA	NCT00876863	NA	
	IT	NA	GAN	Phase I	20	9	Gigaxonin	JeT	NA		NA	NA	NA	O	NCT02362438	NA
		Lom	CLN6	Phase I/II	6	9	CLN6	CAG	$1.5 \times 10^{13}$ vg/kg		NA	NA	NA	O	NCT02725580	NA
IV	PeV	SMA I	Phase I/II	15	9	SMN	CAG	$6.7 \times 10^{13}$ – $3.3 \times 10^{14}$ vg/kg		NA	NA	NA	O	NCT02122952	NA	
	PeV	MPS IIIA	Phase I/II	9	9	SGSH	U1a	$5 \times 10^{12}$ – $1 \times 10^{13}$ vg/kg		NA	NA	Y	O	NCT02716246	NA	

<sup>a</sup>The CAG promoter designation includes the CBA and CB promoter.

Alz, Alzheimer; C, completed; Can, Canavan; Cer, cerebellum; CLN6, Batten CLN6 disease; GAN, giant axonal neuropathy; IS, immunosuppressor; IT, intrathecal; IV, intravenous; LINCF, Late infantile neuronal ceroid lipofuscinosis; MLD, metachromatic leukodystrophy; max vg, maximum vector genome; min vg, minimum vector genome; MPS IIIA, mucopolysaccharidosis IIIA; MPS IIIB, mucopolysaccharidosis IIIB; Lom, lumbar; N, no; NA, not available; NBM, nucleus basalis of meynert; O, ongoing; Par, Parkinson; PeV, peripheral vein; Put, Putamen; SMA I, spinal muscular atrophy type I; SN, substantia nigra; StN, subthalamic nucleus; Str, striatum; WM, white matter; Y, yes.

however, transduction of astrocytes, oligodendrocytes, or microglia is limited<sup>5,6,9</sup> but can be improved by the use of cell-specific promoters.<sup>10–13</sup> Transduction efficiency of these various serotypes has been further established in large-animal models such as dogs, cats, primates, and pigs (Tables 2–4), and finally used in therapeutic trials in human patients (Table 1).

Several strategies have been developed to deliver genes to the CNS.<sup>14</sup> Initially AAV gene delivery to the brain relied on intraparenchymal injection(s) into brain regions affected by the considered disease (Fig. 1). This is a straightforward strategy for conditions involving a restricted brain region, such as, Parkinson's disease. This approach has also been evaluated in conditions affecting large regions of the brain such as lysosomal storage diseases (LSDs), with major successes in terms of preclinical proof of efficacy in numerous animal models (Table 2) and also encouraging results, with excellent safety, in human patients (Table 1) using multiple injection tracks into the brain parenchyma.<sup>15–17</sup>

Many research groups are developing cerebrospinal fluid (CSF)-based delivery, using intracerebroventricular (ICV) or cisternal or lumbar intrathecal (IT) administration (Fig. 1). These ap-

proaches require high quantities of vector compared with intraparenchymal injections (Tables 1–4) but are able to deliver secreted proteins, such as growth factors, lysosomal enzymes, or apolipoprotein E (APOE). However, these strategies are associated with increased leakage of the recombinant vector into the blood circulation and with consequent off-target tissue transduction and expression in peripheral organs, particularly the liver.<sup>18,19</sup> Moreover, strategies will be required to reduce an immune response that may ultimately impair therapeutic benefit in patients.<sup>20</sup>

Several studies have focused on evaluating the ability of AAV vectors to cross the BBB in mouse models and nonhuman primates (NPHs) after IV administration.<sup>21–23</sup> Even if encouraging results could be obtained in neonate mice, decreased transduction efficiency was observed in the primate brain, with a shift in rAAV9 tropism from neuronal to glial cells<sup>23</sup> throughout the brain and, interestingly, into the dorsal root ganglia neurons and motor neurons within the spinal cord.<sup>21</sup> A study comparing the ability of various rAAV strains to cross the BBB in mice demonstrated that rAAVrh.10 is at least as efficient as AAV9 vectors in CNS gene transfer after systemic administration.<sup>24</sup>

**Table 2. Preclinical trials for lysosomal storage diseases**

	Model	Injection site	Disease	Serotype	Transgene	Promoter <sup>a</sup>	Dose, min vg	Dose, max vg	Volume, $\mu$ L	Speed, $\mu$ L/min	Ref.
Intraperitoneal	Rodent	Mice	Tha/DCN or Hip	GM1	1	BGAL	CAG	$4 \times 10^{10}$ – $7.2 \times 10^{10}$	4/1	0.2/0.1	38,166
			Str	GM2	2	HEX	NA	$9.9 \times 10^9$ – $1.4 \times 10^{10}$	3	NA	167
			Cor/Hip/Tha	GCL <sup>b</sup>	2/5	GALC	CAG	$3.3 \times 10^7$	12	NA	168
			Cor/Hth/Cer	INCL <sup>b</sup>	5	CLN1	CAG	$1.2 \times 10^{10}$	12	NA	169
			Str/Hip/Cer	JNCL <sup>b</sup>	rh10	CLN3	CAG	$3 \times 10^{10}$	0.5	1	170
			Tha/Cer	LINCL	2 or 5	CLN2	CAG	$3.6 \times 10^9$	18	0.5	171
			Str	MPS I	2 or 5	IDUA	PGK	$1 \times 10^9$	5	0.5	36
			Str	MPS IIIA	rh10	SGSH/SUMF1	PGK	$7.5 \times 10^9$	2.5	0.5	172
			Str	MPS IIIB	2 or 5	NAGLU	PGK	$1 \times 10^9$	5	0.5	35
			Cor/Cer	MPS IIIB <sup>b</sup>	5	NAGLU	CAG	$1.8 \times 10^{10}$	12	NA	173,174
			Str	MPS VII	5	GUSB	NA	$2.4 \times 10^9$	5	0.5	175
			Cor/Hip/Str	MPS VII	1; 9 or rh10	GUSB	GUSB	$1.2 \times 10^{10}$ – $1.3 \times 10^{10}$	1	0.5	176
			Str or Cor	MLD	rh10	ARSA	CAG	$2.3 \times 10^9$ – $5 \times 10^{10}$	1/2	0.2	177
			Hip/Hth/Cor/Str/Cer	NPD	2	ASM	CMV	$1.2 \times 10^{11}$	24	0.5	178
	Large animal	Cat	Tha/DCN	GM1	1 or rh8	BGAL	CAG	$3 \times 10^{11}$ – $1.2 \times 10^{13}$	94	2	39
			Tha	GM2	1 or rh8	HEX A	CAG	$3 \times 10^{11}$ – $4.2 \times 10^{12}$	70	2	179
		Dog	Str/WM	MPS I	5	IDUA	PGK	$5 \times 10^{11}$ – $2.1 \times 10^{12}$	320	2	37
			Str/WM	MPS IIIB	5	NAGLU	PGK	$5 \times 10^{11}$ – $2.1 \times 10^{12}$	320	2	37
		NHP	WM or Str/Tha	MLD <sup>c</sup>	5	ARSA	PGK	$3.8 \times 10^{11}$ – $1.9 \times 10^{12}$	120	3	180
			WM	MLD <sup>c</sup>	rh10	ARSA	CAG	$2.2 \times 10^{11}$ – $1.1 \times 10^{12}$	360	0.5	181
WM			LINCL <sup>c</sup>	rh10	CLN2	CAG	$1.8 \times 10^{12}$ – $1.8 \times 10^{12}$	180	1	182	
Str	MPS I <sup>c</sup>	1; 2 or 5	IDUA	PGK	$1.4 \times 10^{10}$	100	NA	183			
Intra-CSF	Rodent	Mice	ICV	MPS I <sup>b</sup>	8	IDUA	CAG	$2 \times 10^{10}$	5	1	184
			IT	MPS I	2	IDUA	CMV	$2 \times 10^9$ – $4 \times 10^{10}$	50–100	>5 min	55
			CM	MPS IIIA	9	SGSH	CAG	$5 \times 10^9$ – $5 \times 10^{10}$	5	NA	49
			ICV	MPS IIIA <sup>b</sup>	5	SGSH/SUMF1	CMV	$1.2 \times 10^{10}$ – $6 \times 10^{10}$	2	NA	45
			CM	MPS IIIB	9	NAGLU	CAG	$3 \times 10^{10}$	NA	NA	50
			CM	MPS IIIB	2	NAGLU	CMV	$1 \times 10^{10}$ – $5 \times 10^{10}$	15	<30 sec	185
			IT	MPS VII <sup>b</sup>	2	GUSB	CMV	$1.5 \times 10^{11}$ – $5 \times 10^{11}$	30–100	NA	54
			ICV	MPS VII	4	GUSB	RSV	$1 \times 10^{10}$	10	0.5	46
			ICV	MPS VII <sup>b</sup>	1; 2 or 5	GUSB	GUSB	$1.8 \times 10^{10}$	4	NA	42
			ICV	GCL <sup>b</sup>	1	GALC	CMV	$6 \times 10^{10}$	4	NA	44
			ICV	GM1 <sup>b</sup>	1	BGAL	CAG	$3.3 \times 10^{11}$	4	NA	43
			ICV	MLD	9 or 1	ARSA	CAG	$1.1 \times 10^{11}$ – $2.3 \times 10^{11}$	20	>10 min	186
			ICV	MLD	1	ARSA	NA	$2 \times 10^{11}$	20	>10 min	187
			ICV	MSD <sup>b</sup>	4 or 9	SUMF1	CMV	$1.2 \times 10^{10}$	6	NA	188
	Large animal	Cat	CM	AMD	1 or 9	MANB	GUSB	$1 \times 10^{13}$	NA	NA	51
			CM	MPS I	9	IDUA	CMV/CAG	$1 \times 10^{12}$ vg/kg	1 mL	NA	53
		Dog	CM	MPS I	9	IDUA	CAG	$1 \times 10^{12}$ vg/kg	1–2 mL	NA	156
			CM	MPS IIIA <sup>c</sup>	9	SGSH	CAG	$2 \times 10^{13}$	1 mL	<10 min	49
			CM	MPS IIIB <sup>c</sup>	9	NAGLU	CAG	$6.5 \times 10^{12}$	NA	NA	50
			CM	MPS VII	9 or rh10	GUSB	CAG	$5 \times 10^{12}$ vg/kg	1–2 mL	>1 min	52
ICV		LINCL	1; 2; 4; 5; 8 or 9	CLN2	CAG	$2 \times 10^{12}$	NA	>25 min	47		
NHP	CM	MPS I <sup>c</sup>	9	IDUA	CAG	$3 \times 10^{12}$ vg/kg	1–2 mL	NA	156		
Intravenous	Rodent	Mice	TaV	GM1	9	BGAL	CAG	$1 \times 10^{11}$ – $3.0 \times 10^{13}$	200	NA	61
			TaV	GM2	9	HEX	NA	$3.5 \times 10^{13}$ vg/Kg	100	NA	62
			TaV	GCL <sup>b</sup>	rh10	GALC	CAG	$7.6 \times 10^9$ – $2 \times 10^{11}$	25/10	NA	58,189
			TeV	GCL <sup>b</sup>	2/5	GALC	CAG	$1.38 \times 10^7$	100	NA	168
			JuV	MLD <sup>b</sup>	9	ARSA	CAG	$2 \times 10^{12}$	100	NA	57
			TeV	MSD <sup>b</sup>	4 or 9	SUMF1	CMV	$2 \times 10^{11}$	100	NA	188
			TeV	MPS II <sup>b</sup>	5	IDS	CMV	$1 \times 10^{11}$	NA	NA	190
			TaV	MPS II	8	IDS	EF1 $\alpha$	$1 \times 10^{11}$	200	NA	191
			TaV	MPS IIIA	rh74	SGSH	U1a	$5 \times 10^{12}$ vg/Kg	150–200	NA	192
			NA	MPS IIIA	9	SGSH	CAG	$1 \times 10^{12}$	200	NA	60
			TaV	MPS IIIB	9	NAGLU	CMV	$1 \times 10^{13}$ vg/Kg	150–200	NA	59
			TaV	MPS IIIB	2	NAGLU	CMV	$4 \times 10^{11}$	100–150	NA	193
			NA	MPS VII	9	GUSB	NA	$1 \times 10^{12}$	NA	NA	63
			Large model	Cat	CeV	MPS I	8	IDUA	TBG	$5 \times 10^{12}$ vg/Kg	1 mL
	Dog	CeV		MPS VII <sup>b</sup>	9 or rh10	GUSB	CAG	$2 \times 10^{13}$ vg/Kg	1–2 mL	>2 min	52
	NHP	CeV		MPS IIIB <sup>c</sup>	9	NAGLU	CMV	$1 \times 10^{13}$ – $2 \times 10^{13}$ vg/Kg	5 mL	NA	195

(continued)

Table 2. (Continued)

Model			Injection site	Disease	Serotype	Transgene	Promoter <sup>a</sup>	Dose, min vg	Dose, max vg	Volume, μL	Speed, μL/min	Ref.
IP & CSF	Large	Cat	Tha/ICV	GM2	rh8	HEX	CAG	8 × 10 <sup>11</sup> –1 × 10 <sup>12</sup>		340	2 or 15	196,197
		Rodent	TeV/ICV	MSD <sup>b</sup>	4 or 9	SUMF1	CMV	1.2 × 10 <sup>10</sup> –2 × 10 <sup>11</sup>		103	NA	188
IV CSF	Rodent	Mice	TaV/CM	MPS IIIB	2	NAGLU	CMV	5 × 10 <sup>10</sup> –4 × 10 <sup>11</sup>		5–20	>2 min	198
			TeV/IT	MPS VII <sup>b</sup>	2	GUSB	CMV	3 × 10 <sup>10</sup> –4 × 10 <sup>11</sup>		NA	NA	199
			Large	Dog	CeV/CM	MPS VII <sup>b</sup>	9 or rh10	GUSB	CAG	2.8 × 10 <sup>12</sup> –2 × 10 <sup>13</sup> vg/Kg		1–2 mL

<sup>a</sup>The CAG promoter designation includes the CBA and CB promoters.

<sup>b</sup>Newborn animals.

<sup>c</sup>The study was performed using wild type model of the animal in order to study the adeno-associated virus for the disease mentioned.

AMD, α-mannosidosis; CeV, cephalic vein; CM, cisterna magna; Cor, cortex; DCN, deep cerebellar nuclei; GM, gangliosidosis; GCL, globoid cell leukodystrophy; Hip, hippocampus; Hth, hypothalamus; ICV, intracerebroventricular; INCL, infantile neuronal ceroid lipofuscinosis; JNCL, juvenile neuronal ceroid lipofuscinosis; LINCL, late infantile neuronal ceroid lipofuscinosis; MLD, metachromatic leukodystrophy; MPS, mucopolysaccharidosis; MSD, multiple sulfatase deficiency; NPD, Niemann-Pick disease; IP, intraparenchymal; JuV, jugular vein; Str, striatum; TaV, tail vein; TeV, temporal vein; Tha, thalamus.

## rAAV GENE THERAPY FOR NEURODEGENERATIVE DISEASES

Progress in rAAV gene therapy has allowed for novel treatments of both genetic and acquired neurodegenerative diseases such as LSDs, Alzheimer's disease (AD), Parkinson's disease (PD), Huntington's disease (HD), amyotrophic lateral sclerosis (ALS), and spinal muscular atrophy (SMA). Exciting treatment results have been obtained in animal models of the corresponding human diseases and clinical trials have been, or soon will be, launched.

### Lysosomal storage disorders

LSDs are a group of more than 50 individual inherited metabolic diseases resulting from the deficiency of a lysosomal function responsible for progressive storage of undigested or partially digested materials, leading to cellular and metabolic dysfunction.<sup>25</sup> Although individually rare, they have a collective incidence of approximately 1 per 7000 live births,<sup>26</sup> and at least 70% of all LSDs have a significant CNS component that is typically characterized by neuroinflammation and neurodegeneration in multiple brain regions.<sup>27</sup>

Several features of LSDs make them ideal candidates for gene therapy. Each LSD is a single-gene recessive disorder with a relatively well-understood underlying molecular basis. Cross-correction opens the possibility of genetically modifying a relatively small number of cells so that they express and secrete supraphysiological levels of the deficient enzyme, which can then correct a wide range of cell types at a distance.<sup>28</sup> A portion of newly synthesized lysosomal enzymes can be secreted and internalized by neighboring cells via mannose 6-phosphate receptors that are present on the plasma membrane. In the CNS, circulating lysosomal enzymes can diffuse via the ventricular system or be transported

from the site of production to distal sites via axonal transport.<sup>29</sup> Levels of induced gene expression are generally not critical. Low levels of residual enzyme activity resulting from missense mutations are often associated with significant impacts on the clinical course of the disease.<sup>30</sup> Enzyme activities of less than 10% of the physiological level may therefore be sufficient to have a clinical impact on the disease course. Tight regulation of gene expression is not required because no deleterious effects are associated with supraphysiological levels of most of the lysosomal enzymes. The identification or development of small- and large-animal models recapitulating the clinical course of LSDs has been and is instrumental in evaluating therapeutic strategies.<sup>31</sup>

Several strategies of rAAV infusion have been developed and tested in animal models of neuropathic LSDs via systemic administration as well as direct injection into the CNS through the brain parenchyma or the CSF. Main preclinical studies are listed in Table 2 and results of this research are summarized below.

**Intraparenchymal delivery.** Intraparenchymal injection may effectively treat the CNS of patients with neuropathic LSDs because rAAV and secreted enzymes can be transported to areas distal from the injection sites.<sup>29,32</sup> The relative low dose required, compared with systemic or CSF administration,<sup>33</sup> in an immune-privileged site like the brain also limits the impact of a potential pre-existing immunity to AAV serotypes.<sup>34</sup> Various sites of injection into the brain (cortex, striatum, hippocampus, thalamus, ventral tegment, cerebellum), using various AAV serotypes (1, 2, 5, 9, rh8, and rh10), have been tested in mouse models of neuropathic LSDs such as GM<sub>1</sub> and GM<sub>2</sub> gangliosidosis, globoid cell leukodystrophy (GCL),

**Table 3. Preclinical trials for neurodegenerative diseases**

Model	Injection site	Disease	Serotype	Transgene	Promoter <sup>a</sup>	Dose, min vg	Dose, max vg	Volume, $\mu$ L	Speed, $\mu$ L/min	Ref.		
Intracerebral	Rodent	<b>&gt; Disease affecting the CNS</b>										
		Hip/Cor	Alz	5 or 9	NEP/IDE	NA	NA	NA	NA	NA	68–70	
		Hip/Cor	Alz	5	ECE	CAG	$4 \times 10^9$		4	0.5	69	
		Hip/Cor	Alz	2	ScFv59	CAG	$2 \times 10^9$		8	1	70	
		Cor &/or Hip	Alz	5	CYP46A1	PGK	$2.4 \times 10^9$		4	0.2	87,88	
		Hip	Alz	9	AAPs $\alpha$	Synapsin	$2 \times 10^{10}$		4	0.2	76	
		Hip	Alz	8	IGF-1/IGF-2	EF1 $\alpha$	NA		1	0.2	81	
		Hip	Alz	2/1	IL-10	NA	$3 \times 10^9$		2	NA	82	
		Hip	Alz	1/2	IL-4	NA	$2 \times 10^9$		2	NA	83	
		Hip	Alz	1/2	CCL2 mutant	CAG	$2 \times 10^{10}$		2	NA	200	
		Str/Hip/SN	Par	2 or 5	GBA1/ $\alpha$ -Synuclein	Synapsin	$2 \times 10^{10}$		10	1	93	
		SN	Par	2	Nurr1 &/or Foxa2	CMV	$1 \times 10^9$		1	0.66	94	
		SN	Par	1/2	shRNA anti ROCK2 or LIMK1	Synapsin	$2.5 \times 10^7$ – $1 \times 10^8$		1	0.5	201	
		Str	Par	2	NTN	CAG	$4 \times 10^9$		4	0.2	102	
		Str	Hun	1	DN-TNF	CAG	$2 \times 10^{10}$ – $4.2 \times 10^{10}$		6	0.5	202	
		Str	Hun	1	mRNAs anti-Htt	CAG	$1.8 \times 10^{10}$		4	0.5	117	
		Str	Hun	1	mRNAs anti-Htt	mU6	$4 \times 10^{10}$		10	NA	203	
		Str	Hun	2	NTN	CAG	$4 \times 10^9$		4	2	113	
		Str	Hun	2	CNTF	CAG	$2.7 \times 10^9$		1	0.5	114	
		Str/Cer	Hun	1	shHD2.1	CMV	$6 \times 10^{10}$		12	0.25	116	
	Str	Hun	1/2	ZF11xHunt	CAG	$4.4 \times 10^9$		6	0.25	119		
	Str	Hun	5 or rh10	shCYP46A1	CAG or PU6	$6 \times 10^9$		4	0.2	120		
	Str	Hun	2	GDNF	CAG	$4 \times 10^9$		4	0.2	112		
	Rat	Str	Hun <sup>b</sup>	2	BDNF	CAG	$8 \times 10^8$ – $8 \times 10^9$		5	0.15	109	
	Str	Hun <sup>b</sup>	1/2	BDNF	CAG	$4 \times 10^9$		4	NA	204		
	Str	Hun	2	GDNF	CAG	$1.2 \times 10^{10}$		4	0.5	111		
	Str	Hun <sup>b</sup>	1/2	HD70/HD20/HDh8/shHD2	NSE	$3 \times 10^9$		3	0.1	115		
	Med sep	Alz	NA	NGF	NSE	$6 \times 10^9$		2	NA	205		
	Mice	DCN	ALS	1 or 2	IGF-1	NA	$4 \times 10^{10}$		6	0.5	125	
	Large animal	NPH	<b>&gt; Disease affecting the CNS</b>									
			Put	Hun <sup>b</sup>	1	siRNA	U6	$6.8 \times 10^{10}$		68	1	206
			Cau/Put	Par <sup>b</sup>	2	AADC	NA	$3.6 \times 10^{11}$		180	NA	2
			Put	Par <sup>b</sup>	2	GDNF	CMV	$9.9 \times 10^{11}$		150	NA	100
			Put/SN	Par <sup>b</sup>	2	GDNF	CMV	$8.3 \times 10^{10}$ – $8.3 \times 10^{11}$		50–75	NA	99
			Cau/Put	Par <sup>b</sup>	2	NTN	CAG	$3 \times 10^{11}$		150	2	103
			STN	Par <sup>b</sup>	NA	GAD	NA	$6 \times 10^{10}$ – $1.2 \times 10^{11}$		20	NA	207
			Put	Par <sup>b</sup>	2	hAADC	CMV	$6 \times 10^9$ – $5 \times 10^{11}$		200	0.1–1	101
			Put	Par <sup>b</sup>	2	hAADC-2	CMV	$3 \times 10^{11}$		200	1	97
	Str/SN	Par <sup>b</sup>	NA	GDNF	CAG	$8.4 \times 10^{10}$		21	0.25	98		
	IntraCSF	Rodent	<b>&gt; Disease affecting the CNS</b>									
			ICV	Hun	4	BDNF/ $\Delta$ B2Noggin	NA	$3.7 \times 10^9$		3	NA	110
		Mice	<b>&gt; Disease affecting the spinal cord</b>									
			CVL	ALS	9	amiR SOD1	CAG	$5 \times 10^{10}$		2	NA	208
ICV			ALS	6 or 9	miRNA $\alpha$ SOD1	CMV or GFAP	$2.4 \times 10^{12}$		10	NA	127	
IT			ALS	rh10	amiR-SOD1	Pol II	$1.6 \times 10^{11}$ or $6.8 \times 10^{11}$		3	NA	209	
ICV			ALS	4	IGF-1 or VEGF or IGF-1+VEGF	Hb9	$1.2 \times 10^{12}$ vg/Kg		8	NA	126	
CM			GAN	9	GAN	CMV	$4 \times 10^{10}$ – $8 \times 10^{10}$		20	0.5	210	
ICV			SMA	9	SMN	CAG	$7.2 \times 10^{10}$		10	NA	19	
ICV			SMA	8	SMN	CAG/GUSB	$2.7 \times 10^{12}$ – $3.3 \times 10^{13}$ vg/Kg		5	NA	132	
ICV			SMA	9	SMNI	CAG	$5 \times 10^{10}$ – $1.7 \times 10^{10}$		12	NA	136	
ICV			SMA	9	SMNI	CAG	$1 \times 10^9$ – $5 \times 10^{10}$		6	NA	136	
Large model			Pig	<b>&gt; Disease affecting the spinal cord</b>								
	IT	SMA <sup>b</sup>	9	SMNI	CAG	$3 \times 10^{12}$		1500	NA	136		
NHP	CM/IT	SMA <sup>b</sup>	9	SMNI	CAG	$2.5 \times 10^{13}$		6000	NA	136		
	IT	ALS <sup>b</sup>	rh10	miR SOD1	U6/CAG	$6 \times 10^{12}$ vg/kg		300	NA	211		

(continued)

Table 3. (Continued)

Model			Injection site	Disease	Serotype	Transgene	Promoter <sup>a</sup>	Dose, min vg	Dose, max vg	Volume, $\mu$ L	Speed, $\mu$ L/min	Ref.	
Intravenous	Rodent model	Mice	<b>&gt; Disease affecting the CNS</b>										
			JuV	Hun	9	HTT-specific RNAi	U6		$6.3 \times 10^{11}$ – $3 \times 10^{12}$		50–300	NA	212
			<b>&gt; Disease affecting the spinal cord</b>										
			FaV	SMA	9	SMN	CAG		$5 \times 10^{11}$		NA	NA	131
			FaV	SMA	9	SMN	CMV		$1 \times 10^{11}$		10	NA	133
			TeV	SMA	9	SMN	PGK		$4.5 \times 10^{10}$		70	NA	134
			FaV	SMA	9	SMN	CAG		$5 \times 10^{11}$		100	NA	135
			NA	SMA	9	SMN	CAG		$3.3 \times 10^{14}$ vg/kg		50	NA	19
			TaV	ALS	9	ADAR2	SYN1		$1.5 \times 10^{11}$ – $2.1 \times 10^{12}$		NA	>1 min	130
TaV	ALS	rh10	miR SOD1	U6/CAG		$2 \times 10^{11}$		200	NA	211			
Intramuscular	Rodent	Mice	<b>&gt; Disease affecting the spinal cord</b>										
			Qua	ALS	2	IGF-1 GDNF	NA		$4 \times 10^7$ – $3 \times 10^{10}$		NA	NA	123
Intra SP	Rodent model	Mice	<b>&gt; Disease affecting the spinal cord</b>										
			VGM	ALS	2	IGF-1	CAG		$4.8 \times 10^{10}$		6	0.75	124
			LSC	ALS	2	IL-10	CAG		NA		1	NA	128
	Rat		VH	ALS	2	IGF-1	CAG		$4.1 \times 10^{10}$		5	NA	213

<sup>a</sup>The CAG promoter designation includes the CB and CBA promoter.

<sup>b</sup>The study was performed using wild type model of the animal in order to study the disease mentioned

ALS, amyotrophic lateral sclerosis; Alz, Alzheimer; Cau, caudate; FaV, facial vein; GAN, giant axonal neuropathy; Hun, Huntington; LC, lumbar cord; LSC, lumbar spinal cord; Med Sep, medial septum; Qua, quadriceps; SMA, spinal muscular atrophy; STN, sub-thalamic nucleus; Ven, ventricle; VGM, ventral gray matter; VH, ventral horn.

Niemann-Pick disease (NPD) type A, infantile and late infantile neuronal ceroid lipofuscinosis (INCL and LINCL), metachromatic leukodystrophy (MLD), and mucopolysaccharidosis (MPS) types I, III, and VII (Table 2). These studies showed that AAV vectors, injected into the parenchyma, are able to lead to widespread enzyme distribution as well as biochemical and histological correction in


large regions of the mouse brain, resulting in improved behavioral symptoms, motor function, and life span. These initial data obtained in LSD mice were replicated in large-animal disease models in which multiple parenchymal injections or injection into specific CNS structures with high interconnectivity are more representative of clinical transferability. For example, widespread expres-

Table 4. Preclinical trials for epilepsy and pain

Model			Injection site	Disease	Serotype	Transgene	Promoter <sup>a</sup>	Dose, min vg	Dose, max vg	Volume, $\mu$ L	Speed, $\mu$ L/min	Ref.	
<b>&gt; Epilepsy</b>													
Intraparenchymal	Rodent	Mice	Hip	Epilepsy	8	ADK	gfaABC1D		$2 \times 10^9$		2	1	143
			Hip	Epilepsy	1 and 2	NPY	NSE		$6 \times 10^{10}$		6	0.2	214
			Hip	Epilepsy	1	NPY	CAG		$6.2 \times 10^{10}$		12	1	215
			Hip	Epilepsy	2	GDNF	CAG		$4.1 \times 10^9$ – $6.3 \times 10^9$		3 or 6	0.2	142
			Pir Cor	Epilepsy	2	NPY	CAG		$1.2 \times 10^{10}$ – $1.3 \times 10^{10}$		4	NA	139
			Hip	Epilepsy	1/2 or 2	NPY	NSE		$6.4 \times 10^9$ – $1.3 \times 10^{10}$		6 or 12	NA	138
			Inf Col	Epilepsy	NA	FIB-GAL or GAL	CMV		$1.75 \times 10^7$ – $1 \times 10^9$		1	>9 min	141
			Inf Col or Temp Cor	Epilepsy	NA	NRIA or PINA	CMV		$1 \times 10^9$		1	>9 min	216
			<b>&gt; Neuropathic pain (NP)</b>										
Intra-CSF	Rodent	Mice	IT	Chronic NP	9	shRNA anti TRPV1	hU6		$2 \times 10^{13}$		10	>2 min	148
			IT	Diabetic NP	5	shRNA anti Na <sub>v</sub> 1.3	NA		$3.5 \times 10^{11}$		5	NA	147
			IT	Chronic NP	8	pp $\beta$ EP and/or IL-10	CMV		$3 \times 10^{10}$		15	>1 min	217
Intra SP	Rodent	Rat	DRG	Chronic NP	6	CBD3	CMV		$5 \times 10^9$		2	>5 min	145
			DRG	Chronic NP	2	shRNA anti Na <sub>v</sub> 1.3	hU6		$2.7 \times 10^{10}$ – $1.4 \times 10^{11}$		2	0.4	146
			SGM	Chronic NP	NA	BDNF	CMV		$5 \times 10^5$ IU		1	>2 min	144

<sup>a</sup>The CAG promoter designation includes the CBA promoter.

DRG, dorsal root ganglion; Inf Col, inferior colliculus; NP, neuropathic pain; Pir Cor, piriform cortex; SGM, spinal gray matter; Temp Cor, temporal cortex.



**Figure 1.** Routes of administration targeting the CNS by direct injection into the parenchyma or by injection into the cerebrospinal fluid via the intracerebroventricular or intrathecal (cisternal or lumbar) route.

sion of enzyme throughout the brains of MPS I and MPS IIIB mice after injection of rAAV5 into the striatum was also observed in MPS I and MPS IIIB dogs treated by a similar approach.<sup>35–37</sup> Injection of rAAV1 encoding  $\beta$ -galactosidase into the thalamus and deep cerebellar nuclei of GM<sub>1</sub> gangliosidosis mice was also successful in the feline model, with rAAVrh8 leading to long-term clinical improvement and survival.<sup>38,39</sup> Intraparenchymal injections are particularly applicable to LSDs with severe neurological but moderate systemic involvement, and phase I/II clinical trials have been performed in LINCL, MPS IIIA, MPS IIIB, and MLD (Table 1).

The first phase I/II trial of AAV intraparenchymal injections was for Canavan disease,<sup>40</sup> a pediatric neurodegenerative disorder associated with mutations in the gene encoding aspartoacylase (ASPA),

an enzyme that converts *N*-acetyl-aspartate (NAA) into acetate and aspartate. Results have proved the feasibility and overall safety of rAAV injection into the brain via multiple sites of vector administration. Humoral immune response against the vector was observed in 3 of the 10 subjects.<sup>41</sup> Long-term follow-up indicated a decrease in substrate levels in the brain of some patients and the stabilization of clinical disease.<sup>15</sup> A phase I clinical trial in 10 patients with LINCL was performed by direct injection into the brain parenchyma of rAAV2 carrying the human *CLN2* gene. No safety issue related to the product or the manner of administration was reported, and there was evidence of slowing disease progression.<sup>17</sup> A phase I/II clinical trial in LINCL is currently ongoing with a second-generation AAVrh.10-based vector that confers greater enzyme distribution within the brain and a better immunogenicity profile than rAAV2. rAAVrh.10 was also tested for intracranial delivery of *N*-sulfoglucosamine sulfohydrolase (*SGSH*) and the sulfatase-modifying enzyme factor 1 (*SUMF1*) genes in patients with MPS IIIA. The administration of the vector was safe and well tolerated by the four patients included in the trial after 1 year. Stabilization in three patients and a possible improvement in one patient were observed.<sup>16</sup> In addition, a phase I/II clinical trial for MPS IIIB, using AAV5 to express *N*-acetyl- $\alpha$ -glucosaminidase (NA-GLU), is ongoing as well as a phase I study using AAVrh.10 for MLD (Table 1). Although further follow-up is necessary, these initial results suggest that intraparenchymal AAV-based gene therapy is a realistic option for neuropathic LSDs.

**CSF delivery.** Direct delivery to the CSF is an alternative route that has been tested with various AAV serotypes (1, 2, 4, 5, 8, 9, rh8, and rh10) via ventricles, cisterna magna, or spinal cord into small- and large-animal models of various LSDs such as GM<sub>1</sub> and GM<sub>2</sub> gangliosidosis, GCL, LINCL, MLD, multiple sulfatase deficiency (MSD),  $\alpha$ -mannosidosis (AMD), and MPS I, IIIA, IIIB, and VII (Table 2). Initial proof of concept was made in newborn MPS VII mice where ICV injection of rAAV1, 2, and 5 resulted in the expression of  $\beta$ -glucuronidase (GUSB) in broad regions of the CNS.<sup>42</sup> Similar observations were reported with rAAV1 in newborn mouse models of GCL and GM<sub>1</sub> gangliosidosis<sup>43,44</sup> and with rAAV5 in MPS IIIA newborn pups.<sup>45</sup> Even though broad CNS distribution in newborn mice could be increased by the potential immaturity of the ependymal barrier, or by its disruption favored by the large volume of injection, direct targeting of ependymal cell lining

by rAAV4 in the adult mouse model of MPS VII,<sup>46</sup> or by rAAV2 in the LINCL dog model, also led to broad distribution of enzyme in the CNS associated with therapeutic benefits.<sup>47</sup> However, because of the rapid turnover rate of the ependymal cells,<sup>48</sup> long-term benefits of this approach need to be confirmed. rAAV9 or rAAVrh10 injected into the cisterna magna showed favorable phenotypic outcomes in animal models of MPS types IIIA and B,<sup>49,50</sup> as well as in large-animal models of AMD,<sup>51</sup> MPS VII,<sup>52</sup> and MPS I.<sup>53</sup> Potential additional positive effects of this approach on the non-CNS pathology have been observed but only in the absence of serum antibodies against the vector.<sup>49</sup> Intracisternal administration, although easily accessible in animal models, is more problematic to translate into humans. A less invasive route into the CSF via lumbar intrathecal injection showed positive results of enzyme distribution in the brain of MPS I and MPS VII mouse models, using rAAV2,<sup>54,55</sup> but have not yet been described in the brain of larger animal models. A first clinical trial in Batten CLN6 has been launched with a self-complementary AAV9 carrying the *CLN6* gene administered intrathecally by lumbar puncture (NCT02725580). The minimum calculated dose in this clinical trial (i.e.,  $1.2 \times 10^{14}$  genome copies [GC] for a 1-year-old child of  $\sim 8$  kg) is already 30-fold higher than the highest dose ( $4 \times 10^{12}$  GC) tested so far in intraparenchymal injection clinical trials (Table 1).

**Intravenous delivery.** The capacity of the AAV9 or AAVrh10 serotype to cross the BBB<sup>23,56</sup> has led to the exploration of intravenous (IV) injections as an alternative to treat neuropathic LSD (Table 2). Proof of concept has been shown in neonatal GCL or MLD mice.<sup>57,58</sup> AAV9 IV administration has shown an impact on the CNS pathology of older animal models in MPS III types A and B<sup>59,60</sup> and GM<sub>1</sub> gangliosidosis.<sup>61</sup> However, the efficiency of CNS transduction seems to be more limited in older mice as evaluated in GM<sub>2</sub> gangliosidosis.<sup>62</sup> Moreover, transduction patterns observed in mice can be different from larger animal species.<sup>23</sup> The nature of the storage can also limit the efficacy of this approach, as observed in the MPS VII model where accumulation of sialic acid within the CNS, an inhibitor of AAV9 transduction, precluded any CNS benefit.<sup>63</sup> The first IV phase I/II clinical trials are currently being conducted in spinal muscular atrophy (SMA) (NCT02122952) and MPS IIIA (NCT02716246). Results of the intervention's potential beneficial effect on the CNS aspects of these diseases will demonstrate whether this approach could be envisaged for other LSDs. A significant is-

sue in the transition to humans will be linked to the high doses of virus required to have a therapeutic effect in the CNS (i.e., AAV9 doses up to  $3.3 \times 10^{14}$  GC/kg are proposed in the SMA clinical trial).

Significant progress has been made in global CNS targeting, using rAAV, with evidence of therapeutic potential for neuropathic LSD. Preclinical studies demonstrate that a variety of serotype and injection routes can be used to improve neuropathological and functional deficits. Phase I/II clinical trials of multiple intraparenchymal AAV injections showed the safety of this approach, which can delay the onset or progression of the disease. Studies using larger numbers of patients will be required to confirm these observations.

### Alzheimer's disease

Key neuropathological hallmarks of AD are extracellular amyloid plaques produced from the metabolism of the amyloid precursor protein (APP) and intracellular accumulation of abnormally phosphorylated tau protein, a microtubule assembly protein, forming neurofibrillary tangles (NFTs).<sup>64–66</sup> The mechanisms underlying these neuropathological changes remain unclear. There is today no treatment able to stop the progression of the disease. Several gene therapy strategies have been evaluated, targeting different pathways involved in AD physiopathology, some with encouraging results.

Strategies to reduce the amyloid accumulation by increasing amyloid-degrading proteases, such as neprilysin (NEP) and endothelin-converting enzyme (ECE),<sup>67</sup> or by delivering anti-A $\beta$  single-chain antibody into the corticohippocampal regions of AD mice, using AAV vectors, were shown to decrease plaques and A $\beta$  in the cortex and hippocampus of an AD mouse model<sup>68–70</sup> with no sign of neurotoxicity.<sup>70</sup>

The nonamyloidogenic pathway<sup>71</sup> of APP metabolism prevents the production of amyloid toxic forms and enables the release of the soluble APPs $\alpha$ , which is thought to be responsible for the important physiological functions of APP.<sup>72</sup> Loss of the neuroprotective APPs $\alpha$  could contribute to the development of AD. APPs $\alpha$  levels are decreased in the CSF of patients with AD, in both genetic and sporadic forms, which is correlated with poor memory.<sup>73,74</sup> APPs $\alpha$  inhibits tau phosphorylation through glycogen synthase kinase 3 $\beta$  (GSK3 $\beta$ ) modulation.<sup>75</sup> Increasing the physiological APP pathway is thus an interesting strategy to treat AD, and APPs $\alpha$  overexpression could alleviate AD-related symptoms. APPs $\alpha$  overexpression by means of AAV in the hippocampal neurons of AD mice<sup>76</sup> rescued spatial memory de-

fects, restored synaptic plasticity and spine density, and decreased soluble A $\beta$  and amyloid plaques. This was associated with microglial activation and amyloid plaque phagocytosis.

Neurotrophic factors are good candidates in neuroprotective strategies, but these factors may cause off-target adverse effects, necessitating a targeted delivery strategy to control their localization and spread in the brain.<sup>77</sup> Nerve growth factor (NGF), which stimulates the function of basal forebrain cholinergic neurons undergoing early degeneration in AD,<sup>78</sup> was evaluated in early disease onset in animal models. Encouraging preclinical data and the first clinical trial based on modified fibroblast implantation resulted in a phase I clinical trial in which NGF was administered to 10 patients, who received rAAV2-NGF into the basal forebrain region (NCT00087789). A dose escalation protocol was used:  $1.2 \times 10^{10}$  to  $1.2 \times 10^{11}$  vector particles<sup>79</sup> (Table 1). The brains of patients, which were examined, exhibited a trophic response to NGF in the form of axonal sprouting toward the NGF source. Cholinergic neuronal hypertrophy occurred on the NGF-treated side. Activation of cellular signaling and functional markers was present in the two patients who underwent rAAV2-NGF gene transfer. An overall lower rate of cognitive decline and increased cortical glucose uptake were reported. A phase II multicenter, sham surgery-controlled trial of NGF in AD is ongoing in 49 patients with mild to moderate AD, based on a single administration of AAV-NGF vector that encodes the gene for nerve growth factor (CERE-110) or an appropriate sham (placebo) surgery control treatment.<sup>80</sup> Insulin-like growth factor (IGF)-2, which plays a critical role in memory consolidation in rodents, was administered via an AAV2 vector into the hippocampus of aged wild-type mice and in APP Tg2576 mice. IGF-2 expression enhanced memory, promoted dendritic spine formation in wild-type mice, decreased amyloid levels, and rescued behavioral deficits in AD mice, suggesting that IGF-2 may act as an A $\beta$  scavenger.<sup>81</sup>

Increasing evidence demonstrates the role of inflammation in AD. Antiinflammatory cytokine signaling may play an emerging role as neurotransmitters, neuromodulators, and neurohormones in the brain. Targeting the inflammatory pathway was thus evaluated, using interleukin (IL) delivery. Expression of IL-4 or IL-10 in the brain of AD mice resulted in reduced astro/microgliosis, reduced A $\beta$  deposition, increased neurogenesis, and improved spatial learning.<sup>82,83</sup>

The role of lipid metabolism is increasingly evidenced in AD. APOE, a regulator of lipoprotein metabolism in the CNS that plays several impor-

tant roles such as cholesterol transport, neuroplasticity, and inflammation and A $\beta$  clearance and aggregation<sup>84</sup> is the major risk factor for late-onset AD. A gene transfer approach to bathe the cortex of amyloid plaque-bearing transgenic mice with APOE was used by injecting into the lateral ventricles of AD mice an AAV vector expressing the various human APOE alleles to transduce the ependymal layer. Human APOE proteins diffused into the CSF and interstitial fluid (ISF). Human APOE isoforms affected the concentrations of soluble oligomeric A $\beta$  in the ISF, the pace of A $\beta$  fibrillization and deposition, and the extent of periplaque neurotoxic effects. Increase in soluble A $\beta$ , exacerbation of synaptic loss, and an increased number of dystrophic neurites around each deposit were observed in AD mice receiving APOE4, whereas a relatively protective effect was observed with APOE2. These results suggest that therapeutic approaches aimed at decreasing APOE4 may be beneficial in AD.<sup>85</sup>

In AD, altered cholesterol metabolism seems to play a pivotal role in the formation of amyloid plaques and in tau pathology.<sup>86</sup> The major exportable form of brain cholesterol is 24(S)-hydroxycholesterol (24S-OHC) generated by the neuronal cholesterol 24-hydroxylase enzyme (CYP46A1). Overexpression of CYP46A1 by administering to the brain an AAV vector carrying CYP46A1 in rodent models of AD reduced the number of amyloid plaques and improved spatial memory in amyloid models,<sup>87</sup> and improved cognitive deficits and impaired long-term depression and spine defects in mice with tauopathy,<sup>88</sup> suggesting that CYP46A1 is a relevant target for AD.

All the approaches described have shown some preclinical efficacy in mouse models of AD and raised hopes for clinical applications in human patients. However, the main hurdle of gene therapy for AD is the diffusion of the lesions in the brain. AD impacts a number of anatomical subregions of the brain that are involved in learning and memory. AAV-based gene therapy strategies for AD should thus rely on safe neurosurgical protocols able to target efficiently early-affected regions by means of improved delivery methods and use of serotypes with good diffusion properties (anterograde and retrograde transport). The development of modified AAV vectors able to efficiently cross the BBB should improve such strategies in future.<sup>1,89–92</sup>

### Parkinson's disease

Many neuroprotective treatments have been developed for PD, which is the second most common neurodegenerative disorder. In addition to deep

brain stimulations and cell therapies, gene therapies have emerged as promising alternatives. For example, it has been shown that lysosomal hydrolyase glucocerebrosidase activity is reduced in PD. Increasing it by gene transfer in mouse and rat models of PD showed neuroprotective effects against dopaminergic neuron degeneration.<sup>93</sup> Protection of these dopaminergic neurons from Parkinson degenerative processes has also been shown by AAV-mediated overexpression of the transcription factors *Nurr1* and *Foxa2*.<sup>94</sup> As for other diseases such as epilepsy, the glial cell line-derived neurotrophic factor (GDNF) has been overexpressed in rats and monkeys with lesions of the dopamine system and showed amelioration of lesion-induced behavioral deficits.<sup>95,96</sup> Nonhuman primates are indeed of interest to validate the clinical usefulness of AAV infusion in brain structures affected by PD such as striatum and putamen,<sup>97</sup> and GDNF overexpression in the striatum and substantia nigra of marmoset monkeys showed behavioral and anatomical efficacy.<sup>98</sup> Clinically relevant effects were also obtained in rhesus macaques, where GDNF overexpression promoted restoration of the dopaminergic system in these regions.<sup>99,100</sup> Another strategy often used in gene therapies for PD consists in overexpressing the enzyme aromatic L-amino acid decarboxylase (AADC). Indeed, AADC is in charge of converting L-dopa, the main medication for PD, to dopamine and is decreased in patients with PD. AAV-mediated overexpression of AADC showed 50% improvement in L-dopa responsiveness<sup>101</sup> and persistent results for at least 8 years.<sup>2</sup> CERRE-120, an AAV2 encoding neurturin (NTN), has been shown to protect dopaminergic neurons in preclinical studies.<sup>102,103</sup> Injection of CERRE-120 into the putamen alone or into both the putamen and substantia nigra in more than 100 people with PD (Table 1) showed no adverse effect for up to 5 years.<sup>104</sup> However, efficacy results were modest with no significant improvement in primary outcome measures.<sup>105,106</sup> To date, several clinical trials are ongoing to evaluate the safety and effectiveness of rAAV-GDNF or rAAV-AADC injection (Table 1). Because of the lack of efficacy of several drugs in phase III, such as creatine and ubiquinone, rAAV-gene therapy for PD seems promising and could offer an interesting alternative.

### Huntington's disease

HD is an autosomal dominant neurodegenerative disorder characterized by chorea, dystonia, progressive cognitive deterioration, and psychiatric disturbances evolving to dementia. The disease usually occurs in mid-life, followed by progressive

aggravation of symptoms and death within 10–20 years.<sup>107,108</sup> The genetic cause of the disease is a CAG (cytosine-adenine-guanine) triplet repeat expansion in the huntingtin gene (*HTT*) exceeding approximately 40 copies. Consequent polyglutamine expansion within the amino-terminal region of the HTT protein causes abnormal folding and accumulation of mutant HTT (mHTT) aggregates in cells. Neuroprotective agents have been tested in animal models to counter the toxic cellular effects of mHTT and improve neuronal survival. Brain-derived neurotrophic factor (BDNF), because of its functional interaction with HTT, has been evaluated in animal models of HD. GDNF family ligands (GFLs), that is, glial cell line-derived neurotrophic factor (GDNF), NTN, and ciliary neurotrophic factor (CNTF), have also been shown to support striatal neurons viability *in vitro* and *in vivo*. When administered via rAAV2/1 vector in mice with a quinolinic acid (QA) lesion, BDNF resulted in reduced motor impairment and striatal damage but toxicity was observed with weight loss and seizure activity, showing that lower concentrations of BDNF were necessary for neuroprotection without side effects.<sup>109</sup> Single ICV delivery of BDNF and noggin to rAAV4-BDNF/noggin-treated R6/2 mice resulted in delayed deterioration of motor function and increased survival.<sup>110</sup> rAAV2-GDNF was also administered 2 weeks before a 3-nitropropionic acid (3NP) lesion was induced in rats<sup>111</sup> or in the HD mouse model (N171-82Q mice),<sup>112</sup> resulting in improved performance and reduced neuronal atrophy. NTN, shown to protect striatal projection in excitotoxic models of HD, improved the performance of N171-82Q transgenic HD mice (Rotarod and clasping tests) after rAAV2-NTN (i.e., CERRE-120) injection at 5 weeks of age but did not improve the weight of transgenic animals or their performance in the cognitive radial arm water maze task compared with control groups.<sup>113</sup> CNTF was shown to support striatal neuron viability *in vitro* and *in vivo*. However, a study has shown that long-term expression of CNTF, using the AAV2 vector, increases pathology in the brain of R6/1 transgenic mice.<sup>114</sup> These results may be due to the high dose (rAAV2-CNTF at  $2.7 \times 10^{12}$  vector genomes [VG]/ml, injected unilaterally or bilaterally). Even neurotrophic factors have already been safely used in clinical trials. Altogether these results indicate that caution must be used to establish doses when considering viral delivery of neurotrophic factors to protect against toxic effects of mHTT while avoiding potential side effects.

Direct therapeutic strategies have been developed to target the causative mHTT and reduce the synthesis of mutant protein and potentially pre-

vent cellular damage. RNA interference strategies have shown that decreasing mHTT improves the HD phenotype in mouse and rat models.<sup>112,115–117</sup> Using AAV vectors, various approaches have been developed using mHTT mRNA (single-nucleotide polymorphism)-specific microRNAs (miRNAs) and small hairpin RNAs (shRNAs) to block the formation of the mutant protein, and most recently, allele-specific blockage of transcription. Allele-specific methods selectively silencing mHTT are the most attractive option for HD gene therapy. However, this approach is challenging, because the mutant gene differs from the wild-type allele only by the number of CAG repeats, making both alleles vulnerable. Importantly, some toxic effects of shRNA *in vivo* were suppressed when placed into artificial miRNA expression systems, and miRNA-based approaches may provide more appropriate biological tools for expressing inhibitory RNAs in the brain.<sup>118</sup> Repression of mHTT production was also tested with zinc finger proteins (ZFPs), using long artificial ZFP chains designed to bind longer CAG repeats more effectively than shorter repeats. This reduced chromosomal expression of the mutant gene. *In vivo*, striatal rAAV2/1 delivery in R6/2 mice revealed repression of mHTT in the brain, resulting in protein aggregate reduction and some improvement of motor and behavior performance, establishing a preliminary proof-of-principle for synthetic transcription factor repressors in the brain.<sup>119</sup>

Cholesterol metabolism is impaired in HD, and restoring cholesterol metabolism is thus an interesting therapeutic target. In HD, a decrease in plasma 24S-OHC, the oxysterol produced by brain-specific cholesterol 24-hydroxylase (CYP46A1), follows disease progression proportionally to motor and neuropsychiatric dysfunction and brain atrophy, as indicated by magnetic resonance imaging (MRI), together with lanosterol and lathosterol precursors (markers of cholesterol synthesis). Increased accumulation of cholesterol was observed in striatal neurons together with reduced levels of cholesterol metabolic precursors. CYP46A1 was shown to be decreased in the putamen of patients with HD and R6/2 mice. Moreover, striatal injection of AAVrh10-CYP46A1 vector in R6/2 mice decreased neuronal atrophy, decreased HTT aggregates, and improved motor deficits, as assessed by Rotarod and clasping behavioral tests. It also restored levels of cholesterol and lanosterol and increased levels of desmosterol that were found *in vitro* to protect striatal neurons expressing mHTT from death.<sup>120</sup> These results strongly confirm that restoring cholesterol metabolism through CYP46A1 overexpression is a relevant therapeutic strategy in HD.

### Amyotrophic lateral sclerosis

ALS is a rapidly progressive neurodegenerative disorder affecting motor neurons in the spinal cord, brainstem, and cortex. All voluntary controlled muscles can be affected and most patients with ALS die from respiratory failure 2 to 5 years from the onset of symptoms. Ten to 20% of ALS cases are inherited. The most studied mutations linked to these familial forms are located in the Cu/Zn superoxide dismutase 1 gene (*SOD1*).<sup>121</sup> In *SOD1*-transgenic animals, motor neurons degenerate and the animals die shortly after the onset of symptoms; these animals are often used as human-like ALS models.<sup>122</sup> rAAV encoding IGF-1 was injected into the respiratory and motor limb muscles of *SOD1*<sup>G93A</sup> mice, taking advantage of the vector retrograde transport from presynaptic terminals of projecting neurons to the projecting cell nucleus. Vector was transported up to motor neurons of the spinal cord, which allowed extended survival and a delay in motor decline.<sup>123</sup> Lepore and colleagues used the same strategy but directed the injection into the lumbar spinal cord parenchyma of *SOD1*<sup>G93A</sup> mice. The long-term expression of IGF-1 was followed by delayed disease onset and extended survival but only in male *SOD1*<sup>G93A</sup> mice.<sup>124</sup> The neuroprotective effect of IGF-1 in ALS was associated with decreased glial cell-mediated release of tumor necrosis factor- $\alpha$  and nitric oxide.<sup>125</sup> The same group also achieved promising results with an rAAV injection into the lateral and fourth ventricles of *SOD1*<sup>G93A</sup> mice.<sup>126</sup> More recently, ICV or IT injection of AAV vector encoding microRNA against *SOD1* improved disease outcome in *SOD1*<sup>G93A</sup> mice, with preservation of muscle innervation and neuroprotection.<sup>127</sup> The abundant evidence showing the importance of trophic factors to motor neurons as well as these preclinical results put forward a reliable strategy that could be brought to the clinic. In addition, IL-10 overexpression, using rAAV injection in the spinal cord of *SOD1*<sup>G93A</sup> mice, produces an immune-modulatory effect with longer survival.<sup>128</sup> More than 80% of ALS cases are sporadic and most patients do not carry the *SOD1* mutation. Aizawa and colleagues showed in 2010 a molecular link between reduced adenosine deaminase acting on RNA 2 (ADAR2) activity and TAR DNA-binding protein (TDP-43) pathology, whose loss from the nucleus and positive cytoplasmic inclusions in motor neurons are characteristics of sporadic ALS.<sup>129</sup> Yamashita and colleagues showed that AAV-ADAR2 delivery in motor neurons of ADAR2 knockout mice (AR2) enabled the restoration of ADAR2 activity and also prevented the progression of motor dysfunction and neuronal death.<sup>130</sup>

A few clinical trials using gene therapy in ALS have been started/completed (NCT00748501, NCT02039401, and NCT01041222) but none with rAAV. Despite this, positive results obtained in the above-mentioned preclinical studies may lead to novel promising studies in the clinic.

### Spinal muscular atrophy

SMA is an autosomal recessive neurodegenerative disease leading to infant mortality caused by a mutation in the gene survival of motor neuron (*SMN*). One approach to mitigating the effects of SMN loss of function is exogenous expression of its gene by introducing viral vectors into motor neurons. Several groups provided strong evidence for effective AAV-mediated gene therapy in SMA transgenic mice. AAV9-mediated gene delivery to replace the SMN protein on day 1 after birth in SMA mice resulted in elevated levels of SMN expression and extended survival.<sup>131</sup> rAAV8-SMN was injected bilaterally into the lateral ventricles and the upper lumbar spinal cord of SMA mice, allowing improvements in behavioral tests indicating a functional neuromuscular junction and increased survival from 15 to 50 days.<sup>132</sup> rAAV9 encoding a codon-optimized version of SMN injected into the facial vein on day 1 after birth resulted in a phenotypic correction, a significant increase in survival,<sup>133</sup> complete correction of motor function, and a major increase in survival from 27 to more than 340 days.<sup>134</sup>

As for SMA, gene therapy has been used to rescue the phenotype of SMA with respiratory distress type 1 mice. This disease is caused by a mutation in the *IGHMBP2* gene. rAAV9-IGHMB2 injection restored protein levels, rescued motor function, and increased the life span.<sup>135</sup>

As for ALS or neuropathic pain, intrathecal injection is a promising strategy and proof of concept to transduce motor neurons in juvenile farm pigs and monkeys has already been shown,<sup>136</sup> supporting the use of gene therapy in clinical studies. To date, one clinical trial is actually ongoing to evaluate the safety and efficacy of intravenous delivery of self-complementary rAAV9-SMN as a potential treatment (Table 1).

## rAAV GENE THERAPY FOR OTHER DISORDERS AFFECTING THE NERVOUS SYSTEM

### Epilepsy

Epilepsy is characterized mainly by seizures caused by an imbalance between excitation and inhibition of electrical activities between neurons.

Even if the cause of most cases of epilepsy is unknown, some result from a brain lesion inducing mostly focal seizures, whereas others are associated with genetic defects resulting in generalized seizures. The identification of genetic cases highlighted several ion channels that helped researchers understand and investigate epilepsy, leading to the development of antiepileptic drugs targeting several ion channels. Seizures are often not sufficiently controlled, with current therapeutics, and new therapies are needed.<sup>137</sup> rAAV gene therapy has been used to control seizures in several animal models, based mostly on the overexpression of neuropeptide Y (NPY). NPY, a 36-amino acid member of the pancreatic polypeptide family, is an endogenous modulator of epileptic activity. AAV-mediated gene transfer to overexpress NPY in rats with electroencephalogram (EEG) seizures induced by intrahippocampal or intracerebroventricular kainic acid injection, allowed delayed seizure onset and a reduction in seizures by 50–75%, 8 weeks after rAAV injection.<sup>138</sup> Because NPY activates many receptor subtypes and may result in unwanted side effects, Foti and colleagues developed an rAAV expressing a truncated form of NPY that primarily activates the NPY Y2 receptor and obtained significantly prolonged latency of limbic seizures, 1 week after kainic acid injection.<sup>139</sup> AAV-mediated gene transfer of NPY was also shown to safely improve anticonvulsant activity in kainic acid-treated rats without any glial activation or humoral immune response.<sup>140</sup> Other transgenes have been used to attenuate seizures. Galanin, a neuropeptide with anticonvulsant properties combined with a secretory signal, was overexpressed with beneficial effects in kainic acid-treated rats.<sup>141</sup> GDNF overexpression also leads to suppression of seizures in rat models of temporal lobe epilepsy.<sup>142</sup> Downregulation of adenosine kinase (ADK) in mice via AAV-mediated RNA interference completely abolished spontaneous recurrent seizures in ADK-transgenic mice.<sup>143</sup> Thus antiepileptogenic and antiseizure effects of these gene therapy approaches are promising via a decrease in excitatory signals or an increase in inhibitory signals. To date, no clinical trials with AAV-mediated gene transfer are currently ongoing, but all these preclinical studies support their application in the clinic.

### Neuropathic pain

Even if neuropathic pain is often accompanied by tissue injury, it is a complex state that seems to have no obvious cause. Analgesic agents are recommended as first-line treatments, but analgesia has limitations such as a short half-life, lack of

cellular specificity, and undesired potential off-site effects. Tissue injury, and thus the associated pain, is often chronic and localized and therefore suitable for the use of AAV-mediated gene therapies. A few studies have been carried out with AAV vectors to deliver several gene targets and treat neuropathic pain. In a model of sciatic nerve injury in Wistar rats, AAV vector overexpressing BDNF was injected into the dorsal horn of the spinal cord. One week after injection, rescue in several behavioral tasks was observed such as an alleviation of tactile allodynia and thermal hyperalgesia.<sup>144</sup> Another approach used injection of rAAV expressing the Ca<sup>2+</sup> channel-binding domain 3 (CBD3) into the lumbar dorsal root ganglion (DRG) in rats before the spared nerve injury (SNI). All injected rats presented a significant attenuation of behavioral changes in pain behavior such as hyperalgesia after touch with a pin or sensitivity to acetone stimulation.<sup>145</sup> The knockdown of the voltage-gated sodium channel Nav1.3 via rAAV expressing shRNA was also evaluated in rats with SNI. rAAV injections were performed into the DRG; a partial attenuation of nerve injury-induced mechanical allodynia was observed.<sup>146</sup> Intrathecal injection to treat diabetic neuropathic pain in diabetic Sprague Dawley rats also showed promising results with efficient transduction of DRGs and reduction of nociceptive hyperexcitability and neuropathic tactile allodynia associated with diabetic neuropathic pain.<sup>147</sup> This method is rapid, less invasive than DRG injection, and clinically applicable. Intrathecal injection of AAV vector encoding shRNA against vanilloid receptor 1 (TRPV1) into the subarachnoid space of mice, 3 weeks after SNI surgery, showed significant attenuation of thermal hyperalgesia in response to 50°C heat stimulation observed from 7 days after treatment.<sup>148</sup> These encouraging results should prompt evaluation in human patients with chronic/diabetic neuropathic pain.

## PERSPECTIVE

There is today a tremendous amount of preclinical data demonstrating the relevance and feasibility of AAV-based gene therapy to treat CNS disorders, not only for rare genetic diseases but also potentially for more common severe complex conditions. These have already led to a number of phase I/II clinical applications in human patients with encouraging results that should allow further developments to treat larger numbers of patients. Further steps will need to focus on the development

of new tools improving the safety, efficacy and simplicity of gene delivery to the brain.

A potential immune response to AAV vectors is still a matter of debate. Direct intraparenchymal delivery uses considerably lower doses and is less likely to have an impact on a potential immune response. The development of IT, ICV, and eventually IV injection procedures will need much higher vector doses and will raise the question of immune reactions. After IV delivery, systemic exposure to high doses of AAV vectors can trigger the activation of CD8<sup>+</sup> T cell responses directed against the viral capsid, in a dose-dependent manner, as observed in hemophilia clinical trials.<sup>149</sup> Many approaches are currently being developed to address this crucial issue, immunological modulation in patients, detargeting vectors using specific promoters and miRNA sequences,<sup>150–154</sup> and induction of immune tolerance.<sup>155–157</sup>

Improved efficacy of AAV vectors will benefit from the characterization of new serotypes that progressively emerge from the screening of AAV capsid libraries developed by DNA shuffling of existing AAV capsid genes.<sup>89,158–160</sup> This should allow the design of customized tools, targeted to specific brain regions and for specific cell types. Intravenous delivery remains imperfect by the limited efficacy of AAV vectors to cross the BBB. The development of new tools to transiently increase permeability of the BBB and of new serotypes with higher crossing capacity<sup>90–92</sup> will undoubtedly modify further strategies for safe and efficient CNS gene delivery. However, this will raise a consequent and challenging question concerning the ability to produce sufficient amounts of vectors for future clinical applications in a large number of patients.<sup>161</sup>

## ACKNOWLEDGMENTS

The authors thank the Agence Nationale pour la Recherche (ANR, grant 13-BSV1-022-01), and the Fondation France Alzheimer and the Fondation pour la Recherche Médicale, for their support of research by N.C. on Huntington's disease (ANR) and Alzheimer's disease (FA and FRM). M.A. was supported by a grant from the DIM (Domaine d'Intérêt Majeur) Biotherapy Region Ile de France.

## AUTHOR DISCLOSURE

M.H., S.P., and L.G. are full-time employees of Lysogene. N.C. is a founder and owner of founder equity in BrainVectis Therapeutics. M.A. declares no competing interests.

## REFERENCES

- Choudhury SR, Hudry E, Maguire CA, et al. Viral vectors for therapy of neurologic diseases. *Neuropharmacology* 2016 (in press). DOI: 10.1016/j.neuropharm.2016.02.013
- Hadaczek P, Eberling JL, Pivrotto P, et al. Eight years of clinical improvement in MPTP-lesioned primates after gene therapy with AAV2-hAADC. *Mol Ther* 2010;18:1458–1461.
- Mingozzi F, and High KA. Therapeutic *in vivo* gene transfer for genetic disease using AAV: progress and challenges. *Nat Rev Genet* 2011;12:341–355.
- Rabinowitz JE, Rolling F, Li C, et al. Cross-packaging of a single adeno-associated virus (AAV) type 2 vector genome into multiple AAV serotypes enables transduction with broad specificity. *J Virol* 2002;76:791–801.
- Burger C, Gorbatyuk OS, Velardo MJ, et al. Recombinant AAV viral vectors pseudotyped with viral capsids from serotypes 1, 2, and 5 display differential efficiency and cell tropism after delivery to different regions of the central nervous system. *Mol Ther* 2004;10:302–317.
- Davidson BL, Stein CS, Heth JA, et al. Recombinant adeno-associated virus type 2, 4, and 5 vectors: transduction of variant cell types and regions in the mammalian central nervous system. *Proc Natl Acad Sci U S A* 2000;97:3428–3432.
- Dodiya HB, Bjorklund T, Stansell J III, et al. Differential transduction following basal ganglia administration of distinct pseudotyped AAV capsid serotypes in nonhuman primates. *Mol Ther* 2010;18:579–587.
- Klein RL, Dayton RD, Tatom JB, et al. AAV8, 9, Rh10, Rh43 vector gene transfer in the rat brain: effects of serotype, promoter and purification method. *Mol Ther* 2008;16:89–96.
- Cearley CN, and Wolfe JH. Transduction characteristics of adeno-associated virus vectors expressing cap serotypes 7, 8, 9, and Rh10 in the mouse brain. *Mol Ther* 2006;13:528–537.
- Furman JL, Sama DM, Gant JC, et al. Targeting astrocytes ameliorates neurologic changes in a mouse model of Alzheimer's disease. *J Neurosci* 2012;32:16129–16140.
- Lawlor PA, Bland RJ, Mouravlev A, et al. Efficient gene delivery and selective transduction of glial cells in the mammalian brain by AAV serotypes isolated from nonhuman primates. *Mol Ther* 2009;17:1692–1702.
- von Jonquieres G, Mersmann N, Klugmann CB, et al. Glial promoter selectivity following AAV-delivery to the immature brain. *PLoS One* 2013; 8:e65646.
- Weller ML, Stone IM, Goss A, et al. Selective overexpression of excitatory amino acid transporter 2 (EAAT2) in astrocytes enhances neuroprotection from moderate but not severe hypoxia-ischemia. *Neuroscience* 2008;155:1204–1211.
- Simonato M, Bennett J, Boulis NM, et al. Progress in gene therapy for neurological disorders. *Nat Rev Neurol* 2013;9:277–291.
- Leone P, Shera D, McPhee SW, et al. Long-term follow-up after gene therapy for canavan disease. *Sci Transl Med* 2012;4:165ra163.
- Tardieu M, Zerah M, Husson B, et al. Intracerebral administration of AAV rh.10 carrying human SGSH and SUMF1 cDNAs in children with MPSIIIA disease: results of a phase I/II trial. *Hum Gene Ther* 2014;25:506–516.
- Worgall S, Sondhi D, Hackett NR, et al. Treatment of late infantile neuronal ceroid lipofuscinosis by CNS administration of a serotype 2 adeno-associated virus expressing CLN2 cDNA. *Hum Gene Ther* 2008;19:463–474.
- Gray SJ, Nagabhushan Kalburgi S, McCown TJ, et al. Global CNS gene delivery and evasion of anti-AAV-neutralizing antibodies by intrathecal AAV administration in non-human primates. *Gene Ther* 2013;20:450–459.
- Meyer K, Ferraiuolo L, Schmelzer L, et al. Improving single injection CSF delivery of AAV9-mediated gene therapy for SMA: a dose-response study in mice and nonhuman primates. *Mol Ther* 2015;23:477–487.
- Tse LV, Moller-Tank S, and Asokan A. Strategies to circumvent humoral immunity to adeno-associated viral vectors. *Expert Opin Biol Ther* 2015;15:845–855.
- Bevan AK, Duque S, Foust KD, et al. Systemic gene delivery in large species for targeting spinal cord, brain, and peripheral tissues for pediatric disorders. *Mol Ther* 2011;19:1971–1980.
- Foust KD, Nurre E, Montgomery CL, et al. Intravascular AAV9 preferentially targets neonatal neurons and adult astrocytes. *Nat Biotechnol* 2009;27:59–65.
- Gray SJ, Matagne V, Bachaboina L, et al. Pre-clinical differences of intravascular AAV9 delivery to neurons and glia: a comparative study of adult mice and nonhuman primates. *Mol Ther* 2011;19:1058–1069.
- Zhang H, Yang B, Mu X, et al. Several rAAV vectors efficiently cross the blood-brain barrier and transduce neurons and astrocytes in the neonatal mouse central nervous system. *Mol Ther* 2011;19:1440–1448.
- Cox TM, and Cachon-Gonzalez MB. The cellular pathology of lysosomal diseases. *J Pathol* 2012;226:241–254.
- Fuller M, Meikle PJ, and Hopwood JJ. Epidemiology of lysosomal storage diseases: an overview. In: *Fabry Disease: Perspectives from 5 Years of FOS*. Mehta A, Beck M, Sunder-Plassmann G, eds. Oxford PharmaGenesis, Oxford. 2006; Chapter 2.
- Belletto CM, and Scarpa M. Pathophysiology of neuropathic lysosomal storage disorders. *J Inher Metab Dis* 2010;33:347–362.
- Fratantoni JC, Hall CW, and Neufeld EF. Hurler and Hunter syndromes: mutual correction of the defect in cultured fibroblasts. *Science* 1968;162: 570–572.
- Chen F, Vitry S, Hocquemiller M, et al.  $\alpha$ -L-Iduronidase transport in neurites. *Mol Genet Metab* 2006;87:349–358.
- Leinekugel P, Michel S, Conzelmann E, et al. Quantitative correlation between the residual activity of  $\beta$ -hexosaminidase A and arylsulfatase A and the severity of the resulting lysosomal storage disease. *Hum Genet* 1992;88:513–523.
- Hemsley KM, and Hopwood JJ. Lessons learnt from animal models: pathophysiology of neuropathic lysosomal storage disorders. *J Inher Metab Dis* 2010;33:363–371.
- Salegio EA, Samaranch L, Kells AP, et al. Axonal transport of adeno-associated viral vectors is serotype-dependent. *Gene Ther* 201;20:348–352.
- Rosenberg JB, Sondhi D, Rubin DG, et al. Comparative efficacy and safety of multiple routes of direct CNS administration of adeno-associated virus gene transfer vector serotype rh.10 expressing the human arylsulfatase A cDNA to nonhuman primates. *Hum Gene Ther Clin Dev* 2014;25:164–177.
- Treleaven CM, Tamsett TJ, Bu J, et al. Gene transfer to the CNS is efficacious in immunoprime mice harboring physiologically relevant titers of anti-AAV antibodies. *Mol Ther* 2012; 20:1713–1723.
- Cressant A, Desmaris N, Verot L, et al. Improved behavior and neuropathology in the mouse model of Sanfilippo type IIIB disease after adeno-associated virus-mediated gene transfer in the striatum. *J Neurosci* 2004;24:10229–10239.
- Desmaris N, Verot L, Puech JP, et al. Prevention of neuropathology in the mouse model of Hurler syndrome. *Ann Neurol* 2004;56:68–76.
- Ellinwood NM, Ausseil J, Desmaris N, et al. Safe, efficient, and reproducible gene therapy of the brain in the dog models of Sanfilippo and Hurler syndromes. *Mol Ther* 2011;19:251–259.
- Baek RC, Broekman ML, Leroy SG, et al. AAV-mediated gene delivery in adult GM1-gangliosidosis mice corrects lysosomal storage in CNS and improves survival. *PLoS One* 2010; 5:e13468.
- McCurdy VJ, Johnson AK, Gray-Edwards HL, et al. Sustained normalization of neurological disease after intracranial gene therapy in a feline model. *Sci Transl Med* 2014;6:231ra248.
- Janson C, McPhee S, Bilaniuk L, et al. Clinical protocol: Gene therapy of Canavan disease:

- AAV-2 vector for neurosurgical delivery of aspartoacylase gene (ASPA) to the human brain. *Hum Gene Ther* 2002;13:1391–1412.
41. McPhee SW, Janson CG, Li C, et al. Immune responses to AAV in a phase I study for Canavan disease. *J Gene Med* 2006;8:577–588.
  42. Passini MA, Watson DJ, Vite CH, et al. Intraventricular brain injection of adeno-associated virus type 1 (AAV1) in neonatal mice results in complementary patterns of neuronal transduction to AAV2 and total long-term correction of storage lesions in the brains of  $\beta$ -glucuronidase-deficient mice. *J Virol* 2003;77:7034–7040.
  43. Broekman ML, Baek RC, Comer LA, et al. Complete correction of enzymatic deficiency and neurochemistry in the GM<sub>1</sub>-gangliosidosis mouse brain by neonatal adeno-associated virus-mediated gene delivery. *Mol Ther* 2007;15:30–37.
  44. Rafi MA, Zhi Rao H, Passini MA, et al. AAV-mediated expression of galactocerebrosidase in brain results in attenuated symptoms and extended life span in murine models of globoid cell leukodystrophy. *Mol Ther* 2005;11:734–744.
  45. Fraldi A, Hemsley K, Crawley A, et al. Functional correction of CNS lesions in an MPS-IIIa mouse model by intracerebral AAV-mediated delivery of sulfamidase and SUMF1 genes. *Hum Mol Genet* 2007;16:2693–2702.
  46. Liu G, Martins I, Wemmie JA, et al. Functional correction of CNS phenotypes in a lysosomal storage disease model using adeno-associated virus type 4 vectors. *J Neurosci* 2005;25:9321–9327.
  47. Katz ML, Tecedor L, Chen Y, et al. AAV gene transfer delays disease onset in a TPP1-deficient canine model of the late infantile form of Batten disease. *Sci Transl Med* 2015;7:313ra180.
  48. Chauhan AN, and Lewis PD. A quantitative study of cell proliferation in ependyma and choroid plexus in the postnatal rat brain. *Neuropathol Appl Neurobiol* 1979;5:303–309.
  49. Haurigot V, Marco S, Ribera A, et al. Whole body correction of mucopolysaccharidosis IIIA by intracerebrospinal fluid gene therapy. *J Clin Invest* 2013;123:3254–3271.
  50. Ribera A, Haurigot V, Garcia M, et al. Biochemical, histological and functional correction of mucopolysaccharidosis type IIIB by intracerebrospinal fluid gene therapy. *Hum Mol Genet* 2015;24:2078–2095.
  51. Yoon SY, Bagel JH, O'Donnell PA, et al. Clinical Improvement of  $\alpha$ -mannosidosis cat following a single cisterna magna infusion of AAV1. *Mol Ther* 2016;24:26–33.
  52. Gurda BL, De Guilhem De Lataillade A, Bell P, et al. Evaluation of AAV-mediated gene therapy for central nervous system disease in canine mucopolysaccharidosis VII. *Mol Ther* 2016;24:206–216.
  53. Hinderer C, Bell P, Gurda BL, et al. Intrathecal gene therapy corrects CNS pathology in a feline model of mucopolysaccharidosis I. *Mol Ther* 2014;22:2018–2027.
  54. Elliger SS, Elliger CA, Aguilar CP, et al. Elimination of lysosomal storage in brains of MPS VII mice treated by intrathecal administration of an adeno-associated virus vector. *Gene Ther* 1999;6:1175–1178.
  55. Watson G, Bastacky J, Belichenko P, et al. Intrathecal administration of AAV vectors for the treatment of lysosomal storage in the brains of MPS I mice. *Gene Ther* 2006;13:917–925.
  56. Yang B, Li S, Wang H, et al. Global CNS transduction of adult mice by intravenously delivered rAAVrh.8 and rAAVrh.10 and nonhuman primates by rAAVrh.10. *Mol Ther* 2014;22:1299–1309.
  57. Miyake N, Miyake K, Asakawa N, et al. Long-term correction of biochemical and neurological abnormalities in MLD mice model by neonatal systemic injection of an AAV serotype 9 vector. *Gene Ther* 2014;21:427–433.
  58. Rafi MA, Rao HZ, Luzi P, et al. Intravenous injection of AAVrh10-GALC after the neonatal period in *twitcher* mice results in significant expression in the central and peripheral nervous systems and improvement of clinical features. *Mol Genet Metab* 2015;114:459–466.
  59. Naughton BJ, Duncan FJ, Murrey D, et al. Amyloidosis, synucleinopathy, and prion encephalopathy in a neuropathic lysosomal storage disease: the CNS-biomarker potential of peripheral blood. *PLoS One* 2013;8:e80142.
  60. Ruzo A, Marco S, Garcia M, et al. Correction of pathological accumulation of glycosaminoglycans in CNS and peripheral tissues of MPSIIIA mice through systemic AAV9 gene transfer. *Hum Gene Ther* 2012;23:1237–1246.
  61. Weismann CM, Ferreira J, Keeler AM, et al. Systemic AAV9 gene transfer in adult GM<sub>1</sub> gangliosidosis mice reduces lysosomal storage in CNS and extends lifespan. *Hum Mol Genet* 2015;24:4353–4364.
  62. Walia JS, Altaieb N, Bello A, et al. Long-term correction of Sandhoff disease following intravenous delivery of rAAV9 to mouse neonates. *Mol Ther* 2015;23:414–422.
  63. Chen YH, Clafin K, Geoghegan JC, et al. Sialic acid deposition impairs the utility of AAV9, but not peptide-modified AAVs for brain gene therapy in a mouse model of lysosomal storage disease. *Mol Ther* 2012;20:1393–1399.
  64. Hardy J, and Allsop D. Amyloid deposition as the central event in the aetiology of Alzheimer's disease. *Trends Pharmacol Sci* 1991;12:383–388.
  65. Hardy J, Bogdanovic N, Winblad B, et al. Pathways to Alzheimer's disease. *J Intern Med* 2014;275:296–303.
  66. Selkoe DJ. Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* 2001;81:741–766.
  67. Miners JS, Barua N, Kehoe PG, et al.  $\beta$ -degrading enzymes: potential for treatment of Alzheimer disease. *J Neuropathol Exp Neurol* 2011;70:944–959.
  68. Carty N, Nash KR, Brownlow M, et al. Intracranial injection of AAV expressing NEP but not IDE reduces amyloid pathology in APP+PS1 transgenic mice. *PLoS One* 2013;8:e59626.
  69. Carty NC, Nash K, Lee D, et al. Adeno-associated viral (AAV) serotype 5 vector mediated gene delivery of endothelin-converting enzyme reduces  $A\beta$  deposits in APP + PS1 transgenic mice. *Mol Ther* 2008;16:1580–1586.
  70. Fukuchi K, Tahara K, Kim HD, et al. Anti- $A\beta$  single-chain antibody delivery via adeno-associated virus for treatment of Alzheimer's disease. *Neurobiol Dis* 2006;23:502–511.
  71. Zhang YW, Thompson R, Zhang H, et al. APP processing in Alzheimer's disease. *Mol Brain* 2011;4:3.
  72. Weyer SW, Zagrebelsky M, Herrmann U, et al. Comparative analysis of single and combined APP/APLP knockouts reveals reduced spine density in APP-KO mice that is prevented by APPs $\alpha$  expression. *Acta Neuropathol Commun* 2014;2:36.
  73. Endres K, Fahrenholz F. Regulation of  $\alpha$ -secretase ADAM10 expression and activity. *Exp Brain Res* 2012;217:343–352.
  74. Fahrenholz F.  $\alpha$ -Secretase as a therapeutic target. *Curr Alzheimer Res* 2007;4:412–417.
  75. Deng J, Habib A, Obregon DF, et al. Soluble amyloid precursor protein  $\alpha$  inhibits tau phosphorylation through modulation of GSK3 $\beta$  signaling pathway. *J Neurochem* 2015;135:630–637.
  76. Fol R, Braudeau J, Ludewig S, et al. Viral gene transfer of APPs $\alpha$  rescues synaptic failure in an Alzheimer's disease mouse model. *Acta Neuropathol* 2016;131:247–266.
  77. Malkki H. Alzheimer disease: NGF gene therapy activates neurons in the AD patient brain. *Nat Rev Neurol* 2015;11:548.
  78. Kordower JH, Winn SR, Liu YT, et al. The aged monkey basal forebrain: rescue and sprouting of axotomized basal forebrain neurons after grafts of encapsulated cells secreting human nerve growth factor. *Proc Natl Acad Sci U S A* 1994;91:10898–10902.
  79. Rafii MS, Baumann TL, Bakay RA, et al. A phase 1 study of stereotactic gene delivery of AAV2–NGF for Alzheimer's disease. *Alzheimers Dement* 2014;10:571–581.
  80. Tuszynski MH, Yang JH, Barba D, et al. Nerve growth factor gene therapy: activation of neuronal responses in Alzheimer disease. *JAMA Neurol* 2015;72:1139–1147.
  81. Pascual-Lucas M, Viana da Silva S, Di Scala M, et al. Insulin-like growth factor 2 reverses memory and synaptic deficits in APP transgenic mice. *EMBO Mol Med* 2014;6:1246–1262.
  82. Kiyota T, Ingraham KL, Swan RJ, et al. AAV serotype 2/1-mediated gene delivery of anti-inflammatory interleukin-10 enhances neurogen-

- esis and cognitive function in APP+PS1 mice. *Gene Ther* 2012;19:724–733.
83. Kiyota T, Okuyama S, Swan RJ, et al. CNS expression of anti-inflammatory cytokine interleukin-4 attenuates Alzheimer's disease-like pathogenesis in APP+PS1 bigenic mice. *FASEB J* 2010;24:3093–3102.
  84. Spinney L. Alzheimer's disease: the forgetting gene. *Nature* 2014;510:26–28.
  85. Hudry E, Dashkoff J, Roe AD, et al. Gene transfer of human ApoE isoforms results in differential modulation of amyloid deposition and neurotoxicity in mouse brain. *Sci Transl Med* 2013;5:212ra161.
  86. Ghribi O. Potential mechanisms linking cholesterol to Alzheimer's disease-like pathology in rabbit brain, hippocampal organotypic slices, and skeletal muscle. *J Alzheimers Dis* 2008;15:673–684.
  87. Hudry E, Van Dam D, Kulik W, et al. Adeno-associated virus gene therapy with cholesterol 24-hydroxylase reduces the amyloid pathology before or after the onset of amyloid plaques in mouse models of Alzheimer's disease. *Mol Ther* 2010;18:44–53.
  88. Burlot MA, Braudeau J, Michaelsen-Preusse K, et al. Cholesterol 24-hydroxylase defect is implicated in memory impairments associated with Alzheimer-like tau pathology. *Hum Mol Genet* 2015;24:5965–5976.
  89. Castle MJ, Turunen HT, Vandenbergh LH, et al. Controlling AAV tropism in the nervous system with natural and engineered capsids. *Methods Mol Biol* 2016;1382:133–149.
  90. Deverman BE, Pravdo PL, Simpson BP, et al. Cre-dependent selection yields AAV variants for widespread gene transfer to the adult brain. *Nat Biotechnol* 2016;34:204–209.
  91. Choudhury SR, Fitzpatrick Z, Harris AF, et al. *In vivo* selection yields AAV-B1 capsid for CNS and muscle gene therapy. *Mol Ther* 2016 (in press). DOI: 10.1038/mt.2016.84
  92. Choudhury SR, Harris AF, Cabral DJ, et al. Widespread central nervous system gene transfer and silencing after systemic delivery of novel AAV-AS vector. *Mol Ther* 2016;24:726–735.
  93. Rocha EM, Smith GA, Park E, et al. Glucocerebrosidase gene therapy prevents  $\alpha$ -synucleinopathy of midbrain dopamine neurons. *Neurobiol Dis* 2015;82:495–503.
  94. Oh SM, Chang MY, Song JJ, et al. Combined Nurr1 and Foxa2 roles in the therapy of Parkinson's disease. *EMBO Mol Med* 2015;7:510–525.
  95. Kearns CM, and Gash DM. GDNF protects nigral dopamine neurons against 6-hydroxydopamine *in vivo*. *Brain Res* 1995;672:104–111.
  96. Kirik D, Rosenblad C, Burger C, et al. Parkinson-like neurodegeneration induced by targeted overexpression of  $\alpha$ -synuclein in the nigrostriatal system. *J Neurosci* 2002;22:2780–2791.
  97. Sanftner LM, Sommer JM, Suzuki BM, et al. AAV2-mediated gene delivery to monkey putamen: evaluation of an infusion device and delivery parameters. *Exp Neurol* 2005;194:476–483.
  98. Eslamboli A, Cummings RM, Ridley RM, et al. Recombinant adeno-associated viral vector (rAAV) delivery of GDNF provides protection against 6-OHDA lesion in the common marmoset monkey (*Callithrix jacchus*). *Exp Neurol* 2003;184:536–548.
  99. Johnston LC, Eberling J, Pivrotto P, et al. Clinically relevant effects of convection-enhanced delivery of AAV2-GDNF on the dopaminergic nigrostriatal pathway in aged rhesus monkeys. *Hum Gene Ther* 2009;20:497–510.
  100. Kells AP, Eberling J, Su X, et al. Regeneration of the MPTP-lesioned dopaminergic system after convection-enhanced delivery of AAV2-GDNF. *J Neurosci* 2010;30:9567–9577.
  101. Forsayeth JR, Eberling JL, Sanftner LM, et al. A dose-ranging study of AAV-hAADC therapy in Parkinsonian monkeys. *Mol Ther* 2006;14:571–577.
  102. Gasmi M, Brandon EP, Herzog CD, et al. AAV2-mediated delivery of human neurturin to the rat nigrostriatal system: long-term efficacy and tolerability of CERE-120 for Parkinson's disease. *Neurobiol Dis* 2007;27:67–76.
  103. Herzog CD, Dass B, Holden JE, et al. Striatal delivery of CERE-120, an AAV2 vector encoding human neurturin, enhances activity of the dopaminergic nigrostriatal system in aged monkeys. *Mov Disord* 2007;22:1124–1132.
  104. Marks WJ Jr, Baumann TL, and Bartus RT. Long-term safety of patients with Parkinson's disease receiving rAAV2-neurturin (CERE-120) gene transfer. *Hum Gene Ther* 2016;27:522–527.
  105. Herzog CD, Brown L, Kruegel BR, et al. Enhanced neurotrophic distribution, cell signaling and neuroprotection following substantia nigral versus striatal delivery of AAV2-NRTN (CERE-120). *Neurobiol Dis* 2013;58:38–48.
  106. Marks WJ Jr, Bartus RT, Siffert J, et al. Gene delivery of AAV2-neurturin for Parkinson's disease: a double-blind, randomised, controlled trial. *Lancet Neurol* 2010;9:1164–1172.
  107. Vonsattel JP, and DiFiglia M. Huntington disease. *J Neuropathol Exp Neurol* 1998;57:369–384.
  108. Vonsattel JP, Myers RH, Stevens TJ, et al. Neuropathological classification of Huntington's disease. *J Neuropathol Exp Neurol* 1985;44:559–577.
  109. Kells AP, Henry RA, and Connor B. AAV-BDNF mediated attenuation of quinolinic acid-induced neuropathology and motor function impairment. *Gene Ther* 2008;15:966–977.
  110. Benraiss A, Toner MJ, Xu Q, et al. Sustained mobilization of endogenous neural progenitors delays disease progression in a transgenic model of Huntington's disease. *Cell Stem Cell* 2013;12:787–799.
  111. McBride JL, During MJ, Wu J, et al. Structural and functional neuroprotection in a rat model of Huntington's disease by viral gene transfer of GDNF. *Exp Neurol* 2003;181:213–223.
  112. McBride JL, Ramaswamy S, Gasmi M, et al. Viral delivery of glial cell line-derived neurotrophic factor improves behavior and protects striatal neurons in a mouse model of Huntington's disease. *Proc Natl Acad Sci U S A* 2006;103:9345–9350.
  113. Ramaswamy S, McBride JL, Han I, et al. Intra-striatal CERE-120 (AAV-neurturin) protects striatal and cortical neurons and delays motor deficits in a transgenic mouse model of Huntington's disease. *Neurobiol Dis* 2009;34:40–50.
  114. Denovan-Wright EM, Attis M, Rodriguez-Lebron E, et al. Sustained striatal ciliary neurotrophic factor expression negatively affects behavior and gene expression in normal and R6/1 mice. *J Neurosci Res* 2008;86:1748–1757.
  115. Franich NR, Fitzsimons HL, Fong DM, et al. AAV vector-mediated RNAi of mutant huntingtin expression is neuroprotective in a novel genetic rat model of Huntington's disease. *Mol Ther* 2008;16:947–956.
  116. Harper SQ, Staber PD, He X, et al. RNA interference improves motor and neuropathological abnormalities in a Huntington's disease mouse model. *Proc Natl Acad Sci U S A* 2005;102:5820–5825.
  117. Stanek LM, Sardi SP, Mastis B, et al. Silencing mutant huntingtin by adeno-associated virus-mediated RNA interference ameliorates disease manifestations in the YAC128 mouse model of Huntington's disease. *Hum Gene Ther* 2014;25:461–474.
  118. McBride JL, Boudreau RL, Harper SQ, et al. Artificial miRNAs mitigate shRNA-mediated toxicity in the brain: implications for the therapeutic development of RNAi. *Proc Natl Acad Sci U S A* 2008;105:5868–5873.
  119. Garriga-Canut M, Agustin-Pavon C, Herrmann F, et al. Synthetic zinc finger repressors reduce mutant huntingtin expression in the brain of R6/2 mice. *Proc Natl Acad Sci U S A* 2012;109:E3136–E3145.
  120. Boussicault L, Alves S, Lamaziere A, et al. CYP46A1, the rate-limiting enzyme for cholesterol degradation, is neuroprotective in Huntington's disease. *Brain* 2016;139:953–970.
  121. Kunst CB. Complex genetics of amyotrophic lateral sclerosis. *Am J Hum Genet* 2004;75:933–947.
  122. Higgins CM, Jung C, and Xu Z. ALS-associated mutant SOD1G93A causes mitochondrial vacuolation by expansion of the intermembrane space and by involvement of SOD1 aggregation and peroxisomes. *BMC Neurosci* 2003;4:16.
  123. Kaspar BK, Llado J, Sherkat N, et al. Retrograde viral delivery of IGF-1 prolongs survival in a mouse ALS model. *Science* 2003;301:839–842.

124. Lepore AC, Haenggeli C, Gasmi M, et al. Intraparenchymal spinal cord delivery of adeno-associated virus IGF-1 is protective in the SOD1G93A model of ALS. *Brain Res* 2007;1185:256–265.
125. Dodge JC, Haidet AM, Yang W, et al. Delivery of AAV-IGF-1 to the CNS extends survival in ALS mice through modification of aberrant glial cell activity. *Mol Ther* 2008;16:1056–1064.
126. Dodge JC, Treleaven CM, Fidler JA, et al. AAV4-mediated expression of IGF-1 and VEGF within cellular components of the ventricular system improves survival outcome in familial ALS mice. *Mol Ther* 2010;18:2075–2084.
127. Dirren E, Aebischer J, Rochat C, et al. SOD1 silencing in motoneurons or glia rescues neuromuscular function in ALS mice. *Ann Clin Transl Neurol* 2015;2:167–184.
128. Ayers JI, Fromholt S, Sinyavskaya O, et al. Widespread and efficient transduction of spinal cord and brain following neonatal AAV injection and potential disease modifying effect in ALS mice. *Mol Ther* 2015;23:53–62.
129. Aizawa H, Sawada J, Hideyama T, et al. TDP-43 pathology in sporadic ALS occurs in motor neurons lacking the RNA editing enzyme ADAR2. *Acta Neuropathol* 2010;120:75–84.
130. Yamashita T, Chai HL, Teramoto S, et al. Rescue of amyotrophic lateral sclerosis phenotype in a mouse model by intravenous AAV9-ADAR2 delivery to motor neurons. *EMBO Mol Med* 2013;5:1710–1719.
131. Foust KD, Wang X, McGovern VL, et al. Rescue of the spinal muscular atrophy phenotype in a mouse model by early postnatal delivery of SMN. *Nat Biotechnol* 2010;28:271–274.
132. Passini MA, Bu J, Roskelley EM, et al. CNS-targeted gene therapy improves survival and motor function in a mouse model of spinal muscular atrophy. *J Clin Invest* 2010;120:1253–1264.
133. Valori CF, Ning K, Wyles M, et al. Systemic delivery of scAAV9 expressing SMN prolongs survival in a model of spinal muscular atrophy. *Sci Transl Med* 2010;2:35ra42.
134. Dominguez E, Marais T, Chatauret N, et al. Intravenous scAAV9 delivery of a codon-optimized SMN1 sequence rescues SMA mice. *Hum Mol Genet* 2011;20:681–693.
135. Nizzardo M, Simone C, Rizzo F, et al. Gene therapy rescues disease phenotype in a spinal muscular atrophy with respiratory distress type 1 (SMARD1) mouse model. *Sci Adv* 2015;1:e1500078.
136. Passini MA, Bu J, Richards AM, et al. Translational fidelity of intrathecal delivery of self-complementary AAV9-survival motor neuron 1 for spinal muscular atrophy. *Hum Gene Ther* 2014;25:619–630.
137. Galanopoulou AS, Buckmaster PS, Staley KJ, et al. Identification of new epilepsy treatments: issues in preclinical methodology. *Epilepsia* 2012;53:571–582.
138. Richichi C, Lin EJ, Stefanin D, et al. Anticonvulsant and antiepileptogenic effects mediated by adeno-associated virus vector neuropeptide Y expression in the rat hippocampus. *J Neurosci* 2004;24:3051–3059.
139. Foti S, Haberman RP, Samulski RJ, et al. Adeno-associated virus-mediated expression and constitutive secretion of NPY or NPY13–36 suppresses seizure activity *in vivo*. *Gene Ther* 2007;14:1534–1536.
140. Noe F, Vaghi V, Balducci C, et al. Anticonvulsant effects and behavioural outcomes of rAAV serotype 1 vector-mediated neuropeptide Y overexpression in rat hippocampus. *Gene Ther* 2010;17:643–652.
141. Haberman RP, Samulski RJ, and McCown TJ. Attenuation of seizures and neuronal death by adeno-associated virus vector galanin expression and secretion. *Nat Med* 2003;9:1076–1080.
142. Kanter-Schiffke I, Georgievska B, Kirik D, et al. Seizure suppression by GDNF gene therapy in animal models of epilepsy. *Mol Ther* 2007;15:1106–1113.
143. Theofilas P, Brar S, Stewart KA, et al. Adenosine kinase as a target for therapeutic antisense strategies in epilepsy. *Epilepsia* 2011;52:589–601.
144. Eaton MJ, Blits B, Ruitenber MJ, et al. Amelioration of chronic neuropathic pain after partial nerve injury by adeno-associated viral (AAV) vector-mediated over-expression of BDNF in the rat spinal cord. *Gene Ther* 2002;9:1387–1395.
145. Fischer G, Pan B, Vilceanu D, et al. Sustained relief of neuropathic pain by AAV-targeted expression of CBD3 peptide in rat dorsal root ganglion. *Gene Ther* 2014;21:44–51.
146. Samad OA, Tan AM, Cheng X, et al. Virus-mediated shRNA knockdown of Nav1.3 in rat dorsal root ganglion attenuates nerve injury-induced neuropathic pain. *Mol Ther* 2013;21:49–56.
147. Tan AM, Samad OA, Dib-Hajj SD, et al. Virus-mediated knockdown of Nav1.3 in dorsal root ganglia of STZ-induced diabetic rats alleviates tactile allodynia. *Mol Med* 2015;21:544–552.
148. Hirai T, Enomoto M, Kaburagi H, et al. Intrathecal AAV serotype 9-mediated delivery of shRNA against TRPV1 attenuates thermal hyperalgesia in a mouse model of peripheral nerve injury. *Mol Ther* 2014;22:409–419.
149. Mingozzi F, and High KA. Immune responses to AAV in clinical trials. *Curr Gene Ther* 2011;11:321–330.
150. Borel F, Kay MA, and Mueller C. Recombinant AAV as a platform for translating the therapeutic potential of RNA interference. *Mol Ther* 2014;22:692–701.
151. Geisler A, Jungmann A, Kurreck J, et al. microRNA122-regulated transgene expression increases specificity of cardiac gene transfer upon intravenous delivery of AAV9 vectors. *Gene Ther* 2011;18:199–209.
152. Majowicz A, Maczuga P, Kwikkers KL, et al. Mir-142-3p target sequences reduce transgene-directed immunogenicity following intramuscular adeno-associated virus 1 vector-mediated gene delivery. *J Gene Med* 2013;15:219–232.
153. Qiao C, Yuan Z, Li J, et al. Liver-specific microRNA-122 target sequences incorporated in AAV vectors efficiently inhibits transgene expression in the liver. *Gene Ther* 2011;18:403–410.
154. Xie J, Xie Q, Zhang H, et al. MicroRNA-regulated, systemically delivered rAAV9: a step closer to CNS-restricted transgene expression. *Mol Ther* 2011;19:526–535.
155. Doerfler PA, Todd AG, Clement N, et al. Co-packaged AAV9 vectors promote simultaneous immune tolerance and phenotypic correction of Pompe disease. *Hum Gene Ther* 2016;27:43–59.
156. Hinderer C, Bell P, Louboutin JP, et al. Neonatal systemic AAV induces tolerance to CNS gene therapy in MPS I dogs and nonhuman primates. *Mol Ther* 2015;23:1298–1307.
157. Mingozzi F, and Buning H. Adeno-associated viral vectors at the frontier between tolerance and immunity. *Front Immunol* 2015;6:120.
158. Grimm D, Lee JS, Wang L, et al. *In vitro* and *in vivo* gene therapy vector evolution via multi-species interbreeding and retargeting of adeno-associated viruses. *J Virol* 2008;82:5887–5911.
159. Koerber JT, Jang JH, and Schaffer DV. DNA shuffling of adeno-associated virus yields functionally diverse viral progeny. *Mol Ther* 2008;16:1703–1709.
160. Maheshri N, Koerber JT, Kaspar BK, et al. Directed evolution of adeno-associated virus yields enhanced gene delivery vectors. *Nat Biotechnol* 2006;24:198–204.
161. Grieger JC, and Samulski RJ. Adeno-associated virus vectorology, manufacturing, and clinical applications. *Methods Enzymol* 2012;507:229–254.
162. LeWitt PA, Rezai AR, Leehey MA, et al. AAV2–GAD gene therapy for advanced Parkinson’s disease: a double-blind, sham-surgery controlled, randomised trial. *Lancet Neurol* 2011;10:309–319.
163. Mittermeyer G, Christine CW, Rosenbluth KH, et al. Long-term evaluation of a phase 1 study of AADC gene therapy for Parkinson’s disease. *Hum Gene Ther* 2012;23:377–381.
164. Marks WJ Jr, Ostrem JL, Verhagen L, et al. Safety and tolerability of intraputamin delivery of CER-120 (adeno-associated virus serotype 2-neurturin) to patients with idiopathic Parkinson’s disease: an open-label, phase I trial. *Lancet Neurol* 2008;7:400–408.
165. Bartus RT, Baumann TL, Siffert J, et al. Safety/feasibility of targeting the substantia nigra with AAV2-neurturin in Parkinson patients. *Neurology* 2013;80:1698–1701.
166. Broekman ML, Tierney LA, Benn C, et al. Mechanisms of distribution of mouse  $\beta$ -galactosidase in the adult GM<sub>1</sub>-gangliosidosis brain. *Gene Ther* 2009;16:303–308.

167. Sargeant TJ, Wang S, Bradley J, et al. Adeno-associated virus-mediated expression of  $\beta$ -hexosaminidase prevents neuronal loss in the Sandhoff mouse brain. *Hum Mol Genet* 2011; 20:4371–4380.
168. Lin D, Fantz CR, Levy B, et al. AAV2/5 vector expressing galactocerebrosidase ameliorates CNS disease in the murine model of globoid-cell leukodystrophy more efficiently than AAV2. *Mol Ther* 2005;12:422–430.
169. Macauley SL, Wong AM, Shyng C, et al. An anti-neuroinflammatory that targets dysregulated glia enhances the efficacy of CNS-directed gene therapy in murine infantile neuronal ceroid lipofuscinosis. *J Neurosci* 2014;34:13077–13082.
170. Sondhi D, Scott EC, Chen A, et al. Partial correction of the CNS lysosomal storage defect in a mouse model of juvenile neuronal ceroid lipofuscinosis by neonatal CNS administration of an adeno-associated virus serotype rh.10 vector expressing the human *CLN3* gene. *Hum Gene Ther* 2014;25:223–239.
171. Passini MA, Dodge JC, Bu J, et al. Intracranial delivery of *CLN2* reduces brain pathology in a mouse model of classical late infantile neuronal ceroid lipofuscinosis. *J Neurosci* 2006;26:1334–1342.
172. Winner LK, Beard H, Hassiotis S, et al. A pre-clinical study evaluating AAVrh10-based gene therapy for Sanfilippo syndrome. *Hum Gene Ther* 2016;27:363–375.
173. Heldermon CD, Ohlemiller KK, Herzog ED, et al. Therapeutic efficacy of bone marrow transplant, intracranial AAV-mediated gene therapy, or both in the mouse model of MPS IIIB. *Mol Ther* 2010;18:873–880.
174. Heldermon CD, Qin EY, Ohlemiller KK, et al. Disease correction by combined neonatal intracranial AAV and systemic lentiviral gene therapy in Sanfilippo syndrome type B mice. *Gene Ther* 2013;20:913–921.
175. Liu G, Chen YH, He X, et al. Adeno-associated virus type 5 reduces learning deficits and restores glutamate receptor subunit levels in MPS VII mice CNS. *Mol Ther* 2007;15:242–247.
176. Cearley CN, and Wolfe JH. A single injection of an adeno-associated virus vector into nuclei with divergent connections results in widespread vector distribution in the brain and global correction of a neurogenetic disease. *J Neurosci* 2007;27:9928–9940.
177. Piguet F, Sondhi D, Piraud M, et al. Correction of brain oligodendrocytes by AAVrh.10 intracerebral gene therapy in metachromatic leukodystrophy mice. *Hum Gene Ther* 2012;23:903–914.
178. Passini MA, Bu J, Fidler JA, et al. Combination brain and systemic injections of AAV provide maximal functional and survival benefits in the Niemann-Pick mouse. *Proc Natl Acad Sci U S A* 2007;104:9505–9510.
179. Bradbury AM, Cochran JN, McCurdy VJ, et al. Therapeutic response in feline Sandhoff disease despite immunity to intracranial gene therapy. *Mol Ther* 2013;21:1306–1315.
180. Colle MA, Piguet F, Bertrand L, et al. Efficient intracerebral delivery of AAV5 vector encoding human ARSA in non-human primate. *Hum Mol Genet* 2010;19:147–158.
181. Zerah M, Piguet F, Colle MA, et al. Intracerebral gene therapy using AAVrh.10-hARSA recombinant vector to treat patients with early-onset forms of metachromatic leukodystrophy: pre-clinical feasibility and safety assessments in non-human primates. *Hum Gene Ther Clin Dev* 2015;26:113–124.
182. Sondhi D, Johnson L, De B, et al. Long term expression and safety of administration of AAVrh.10hCLN2 to the brain of rats and non-human primates for the treatment of late infantile neuronal lipofuscinosis. *Hum Gene Ther Methods* 2012;23:324–335.
183. Ciron C, Cressant A, Roux F, et al. Human  $\alpha$ -iduronidase gene transfer mediated by adeno-associated virus types 1, 2, and 5 in the brain of nonhuman primates: vector diffusion and biodistribution. *Hum Gene Ther* 2009;20:350–360.
184. Wolf DA, Lenander AW, Nan Z, et al. Direct gene transfer to the CNS prevents emergence of neurologic disease in a murine model of mucopolysaccharidosis type I. *Neurobiol Dis* 2011; 43:123–133.
185. Fu H, DiRosario J, Kang L, et al. Restoration of central nervous system  $\alpha$ -N-acetylglucosaminidase activity and therapeutic benefits in mucopolysaccharidosis IIIB mice by a single intracisternal recombinant adeno-associated viral type 2 vector delivery. *J Gene Med* 2010;12:624–633.
186. Hironaka K, Yamazaki Y, Hirai Y, et al. Enzyme replacement in the CSF to treat metachromatic leukodystrophy in mouse model using single intracerebroventricular injection of self-complementary AAV1 vector. *Sci Rep* 2015;5: 13104.
187. Yamazaki Y, Hirai Y, Miyake K, et al. Targeted gene transfer into ependymal cells through intraventricular injection of AAV1 vector and long-term enzyme replacement via the CSF. *Sci Rep* 2014;4:5506.
188. Spanpanato C, De Leonibus E, Dama P, et al. Efficacy of a combined intracerebral and systemic gene delivery approach for the treatment of a severe lysosomal storage disorder. *Mol Ther* 2011;19:860–869.
189. Rafi MA, Rao HZ, Luzi P, et al. Extended normal life after AAVrh10-mediated gene therapy in the mouse model of Krabbe disease. *Mol Ther* 2012;20:2031–2042.
190. Polito VA, and Cosma MP. IDS crossing of the blood-brain barrier corrects CNS defects in MPSII mice. *Am J Hum Genet* 2009;85:296–301.
191. Jung SC, Park ES, Choi EN, et al. Characterization of a novel mucopolysaccharidosis type II mouse model and recombinant AAV2/8 vector-mediated gene therapy. *Mol Cells* 2010;30:13–18.
192. Duncan FJ, Naughton BJ, Zaraspe K, et al. Broad functional correction of molecular impairments by systemic delivery of scAAVrh74-hSGSH gene delivery in MPS IIIA mice. *Mol Ther* 2015; 23:638–647.
193. McCarty DM, DiRosario J, Gulaid K, et al. Mannitol-facilitated CNS entry of rAAV2 vector significantly delayed the neurological disease progression in MPS IIIB mice. *Gene Ther* 2009; 16:1340–1352.
194. Hinderer C, Bell P, Gurda BL, et al. Liver-directed gene therapy corrects cardiovascular lesions in feline mucopolysaccharidosis type I. *Proc Natl Acad Sci U S A* 2014;111:14894–14899.
195. Murrey DA, Naughton BJ, Duncan FJ, et al. Feasibility and safety of systemic rAAV9-hNAGLU delivery for treating mucopolysaccharidosis IIIB: toxicology, biodistribution, and immunological assessments in primates. *Hum Gene Ther Clin Dev* 2014;25:72–84.
196. Rockwell HE, McCurdy VJ, Eaton SC, et al. AAV-mediated gene delivery in a feline model of Sandhoff disease corrects lysosomal storage in the central nervous system. *ASN Neuro* 2015;7.
197. Gray-Edwards HL, Brunson BL, Holland M, et al. Mucopolysaccharidosis-like phenotype in feline Sandhoff disease and partial correction after AAV gene therapy. *Mol Genet Metab* 2015;116: 80–87.
198. Fu H, Kang L, Jennings JS, et al. Significantly increased lifespan and improved behavioral performances by rAAV gene delivery in adult mucopolysaccharidosis IIIB mice. *Gene Ther* 2007;14:1065–1077.
199. Elliger SS, Elliger CA, Lang C, et al. Enhanced secretion and uptake of  $\beta$ -glucuronidase improves adeno-associated viral-mediated gene therapy of mucopolysaccharidosis type VII mice. *Mol Ther* 2002;5:617–626.
200. Kiyota T, Yamamoto M, Schroder B, et al. AAV1/2-mediated CNS gene delivery of dominant-negative CCL2 mutant suppresses gliosis,  $\beta$ -amyloidosis, and learning impairment of APP/PS1 mice. *Mol Ther* 2009;17:803–809.
201. Saal KA, Koch JC, Tatenhorst L, et al. AAV-shRNA-mediated downregulation of ROCK2 attenuates degeneration of dopaminergic neurons in toxin-induced models of Parkinson's disease *in vitro* and *in vivo*. *Neurobiol Dis* 2015;73:150–162.
202. Alto LT, Chen X, Ruhn KA, et al. AAV-dominant negative tumor necrosis factor (DN-TNF) gene transfer to the striatum does not rescue medium spiny neurons in the YAC128 mouse model of Huntington's disease. *PLoS One* 2014;9:e96544.
203. Boudreau RL, McBride JL, Martins I, et al. Nonallele-specific silencing of mutant and wild-type huntingtin demonstrates therapeutic efficacy in Huntington's disease mice. *Mol Ther* 2009;17:1053–1063.

204. Connor B, Sun Y, von Hieber D, et al. AAV1/2-mediated BDNF gene therapy in a transgenic rat model of Huntington's disease. *Gene Ther* 2016;23:283–295.
205. Klein RL, Hirko AC, Meyers CA, et al. NGF gene transfer to intrinsic basal forebrain neurons increases cholinergic cell size and protects from age-related, spatial memory deficits in middle-aged rats. *Brain Res* 2000;875:144–151.
206. McBride JL, Pitzer MR, Boudreau RL, et al. Preclinical safety of RNAi-mediated HTT suppression in the rhesus macaque as a potential therapy for Huntington's disease. *Mol Ther* 2011;19:2152–2162.
207. Emborg ME, Carbon M, Holden JE, et al. Subthalamic glutamic acid decarboxylase gene therapy: changes in motor function and cortical metabolism. *J Cereb Blood Flow Metab* 2007;27:501–509.
208. Stoica L, Todeasa SH, Cabrera GT, et al. Adeno-associated virus-delivered artificial microRNA extends survival and delays paralysis in an amyotrophic lateral sclerosis mouse model. *Ann Neurol* 2016;79:687–700.
209. Wang H, Yang B, Qiu L, et al. Widespread spinal cord transduction by intrathecal injection of rAAV delivers efficacious RNAi therapy for amyotrophic lateral sclerosis. *Hum Mol Genet* 2014;23:668–681.
210. Mussche S, Devreese B, Nagabhushan Kalburgi S, et al. Restoration of cytoskeleton homeostasis after gigaxonin gene transfer for giant axonal neuropathy. *Hum Gene Ther* 2013;24:209–219.
211. Borel F, Gernoux G, Cardozo B, et al. Therapeutic rAAVrh10 mediated SOD1 silencing in adult SOD1<sup>G93A</sup> mice and nonhuman primates. *Hum Gene Ther* 2016;27:19–31.
212. Dufour BD, Smith CA, Clark RL, et al. Intrajugular vein delivery of AAV9-RNAi prevents neuropathological changes and weight loss in Huntington's disease mice. *Mol Ther* 2014;22:797–810.
213. Franz CK, Federici T, Yang J, et al. Intraspinal cord delivery of IGF-I mediated by adeno-associated virus 2 is neuroprotective in a rat model of familial ALS. *Neurobiol Dis* 2009;33:473–481.
214. Sorensen AT, Nikitidou L, Ledri M, et al. Hippocampal NPY gene transfer attenuates seizures without affecting epilepsy-induced impairment of LTP. *Exp Neurol* 2009;215:328–333.
215. Noe F, Pool AH, Nissinen J, et al. Neuropeptide Y gene therapy decreases chronic spontaneous seizures in a rat model of temporal lobe epilepsy. *Brain* 2008;131:1506–1515.
216. Haberman R, Criswell H, Snowdy S, et al. Therapeutic liabilities of *in vivo* viral vector tropism: adeno-associated virus vectors, NMDAR1 antisense, and focal seizure sensitivity. *Mol Ther* 2002;6:495–500.
217. Storek B, Reinhardt M, Wang C, et al. Sensory neuron targeting by self-complementary AAV8 via lumbar puncture for chronic pain. *Proc Natl Acad Sci U S A* 2008;105:1055–1060.

Received for publication June 7, 2016;  
accepted after revision June 7, 2016.

Published online: June 7, 2016.

**This article has been cited by:**

1. Roeland Buckinx, Jean-Pierre Timmermans. 2016. Targeting the gastrointestinal tract with viral vectors: state of the art and possible applications in research and therapy. *Histochemistry and Cell Biology* **146**:6, 709-720. [[CrossRef](#)]