

RESEARCH ARTICLE**A Cure for Sanfilippo Syndrome? A Summary of Current Therapeutic Approaches and their Promise****Authors**Yewande Pearse¹ and Michelina Iacovino¹.**Affiliation**¹Department of Pediatrics, The Lundquist Institute at Harbor-UCLA Medical Center, Torrance, CA 90502**Correspondence**Yewande Pearse¹Email: yewande.pearse@lundquist.org

* These Authors contributed equally to this review.

Abstract

Mucopolysaccharidoses III (MPS III, Sanfilippo syndrome) is a subtype of the Mucopolysaccharidoses (MPS), a group of inherited lysosomal disorders caused by a deficiency of lysosomal enzymes responsible for catabolizing glycosaminoglycans (GAGs). Although MPS III is rare, MPS diseases as a group are relatively frequent with an overall incidence of approximately 1 in 20,000 – 25,000 births. MPS III are paediatric diseases, which cause learning difficulties, behavioural disorders and dementia, as well as skeletal deformities and ultimately result in premature death. There are currently no approved treatments for MPS III, but a number of therapeutic approaches are under development. In the past 30 years, research using cellular and animal models have led to clinical trials involving enzyme replacement therapy (ERT), substrate reduction therapy (SRT) and gene therapy, while stem cells approaches remain at the pre-clinical stage. Although safety and clinical efficacy in animal models have shown promise, the results of clinical trials have proved costly and shown limited therapeutic effects. In this review, we describe the most recent results from clinical trials. While ERT and gene therapy are the most developed therapies for MPS III, we highlight the work that needs to be done to bring us closer to a real treatment for these devastating diseases.

Key words: Mucopolysaccharidoses III, Sanfilippo syndrome, lysosomal storage disease, neurodegeneration, enzyme replacement therapy, gene therapy, substrate reduction therapy, stem cell therapy, clinical trial

1.Introduction

Mucopolysaccharidosis type III (MPS III or Sanfilipo Syndrome) refers to one of five (MPS IIIA-E) autosomal recessive lysosomal storage diseases. Each form of MPS III is caused by a mutation in both alleles of a gene which codes for an enzyme involved in the degradation of the glycosaminoglycan (GAG) heparin sulfate (HS). As a result, partially degraded HS accumulates in lysosomes leading to lysosomal malfunction and disease¹⁻³. The subtypes of MPS III are caused by deficiencies in the enzymes; sulfaminase (MPS IIIA, OMIM no. 252900)⁴, α -*N*-acetylglucosaminidase (NAGLU, MPS IIIB, OMIM no. 252920)⁵, heparin acetyl CoA: α -glucosaminide *N*-acetyltransferase (HGSNAT, MPS IIIC, OMIM 252930)⁶, *N*-acetylglucosamine 6-sulfatase (GNS, MPS IIID, OMIM 252940)⁴; and *N*-glucosamine 3-*O*-sulfatase (arylsulfatase G or ARSG, MPS IIIE)⁷. MPS III is the most common form of MPS, with a prevalence of between approximately 0.3 and 4.1 cases for every 100, 000 births depending on the subtype⁸. MPS III manifests in a similar way regardless of subtype, although the age of onset and rate of disease progression may differ between individuals. The most striking feature of MPS III, compared to other forms of MPS is that it primarily affects the brain, distinguishing them as neurological diseases. Although other forms of MPS are linked to severe somatic symptoms and may also have neurological characteristics, only MPS III appears to heavily involve the central nervous system. Pre-natal and early stages of post-natal development appear to be normal in MPS III, with symptoms occurring during the first few

years of life (1-3 years of age) and including developmental delay such as difficulty forming language, cognitive decline, hyperactivity, sleep disturbances, aggressive behaviour and seizures, especially in older children. Towards the later stages of the disease, hyperactivity and anxiety subside but patients succumb to motor impairment, eventually reaching a vegetative state, become less responsive to external stimuli and die prematurely before their third decade of life⁹⁻¹¹. Unsurprisingly, these symptoms have a tremendous impact not just on the children with MPS III, but parents and primary carers of children with MPS III. Currently, there are no available treatments to effectively reverse or slow down disease progression in MPS III. Instead, most efforts are palliative, focusing more on regulating behaviour and sleep disturbances. This is because the clinical symptoms that characterize MPS III result from neuronal dysfunction, making them particularly difficult to treat. MPS types I, II, IVA, and VI for which clinical symptoms are predominantly somatic, can all be treated using Enzyme Replacement Therapy (ERT) or bone marrow and hematopoietic stem cell (HSCT) transplantation¹². Attempts have been made to deliver missing enzyme to MPS III patients via the blood stream, but the results have been poor due to the inability of the enzyme to cross the blood-brain-barrier. Notwithstanding, a number of promising therapies are currently being tested in cell and animal models and several clinical trials are underway. In this review, we will focus on the progress that has been made for ERT, small molecules and stem cell approaches in the

past few years, which obstacles still remain and how they may be overcome.

2. Therapeutic Scope

There are two broad ways in which monogenetic diseases like MPS III can be treated. One, targeting the gene defect directly through gene therapy or two, targeting the deficiency of the protein that the gene codes for using modified ERT, the use of small molecules or stem cells. All four treatment approaches have been tested in cellular and animal models, and clinical trial. An excellent review has been written by Gaffke *et al.*, recently ¹³. Building on these reported findings, this review will concentrate on the work that has been done toward obtaining therapeutic options for MPS III in the last two years and their promise.

2.1. Cellular Models.

Cellular models serve as a useful tool for testing the molecular mechanisms, which drive MPS III, or the efficacy of various therapies. For obvious reasons, cellular models can only provide a preliminary indication of the effects of compounds on an organism. Nonetheless, ERT, substrate reduction therapy and other forms of small molecule therapies have offered promising results, which warrant further investigation.

The advent of three-dimensional (3D) organoid culture systems, generated from induced pluripotent stem cells (iPSCs) provide unprecedented potential for modelling the human brain, mimicking

various developmental features at the molecular and cellular level ¹⁴⁻¹⁹. These scaled-down complex cellular models not only offer a way to better study the mechanism of disease in MPS III, but in addition, better test small molecules. For the first time, two stem cell lines derived from the skin of a patient with MPS IIIA and MPS IIIB, have been generated and characterized ^{20,21}. Although iPSC-derived organoids are yet to be generated from these lines, they represent the first step towards MPS III brain organoids. Much progress has been made using 2D-culture systems as described above, but the greater complexity achieved using 3D-organoids may bridge the gap between *in vitro* and clinical studies.

2.2 Animal Models

Animal models of genetic diseases provide a biological system to test potential therapies that have been previously investigated *in vitro*. The therapies described above have all proved relatively efficacious in cell models. However, preclinical studies in both cell (ideally human) and whole organisms are necessary in order to validate potential therapies with real clinical application. This is particularly important for MPS III, which unlike most forms of MPS, displays primarily neurological symptoms. Therefore, testing these therapies in animal models addresses the issue of a blood-brain-barrier, a structure conserved in both rodents for example, and humans.

2.3. Clinical Trials

Cellular and animal models have proved useful in identifying potential therapeutic

strategies to reduce the primary cause of the disease, GAG accumulation in the brain and its downstream effects. These studies have paved the way for 26 clinical trials for MPS III in the past 20 years, 9 of which have been completed. These include stem cell transplantation, ERT, substrate reduction therapy and gene therapy. One of the main limitations of assessing the effects of treatments in MPS III is that it is a rare disease, meaning that sample sizes tend to be small and fail to completely represent the different stage and severity of symptoms. To overcome this issue, future studies should take heed of recent recommendations on clinical trial design for the treatment of MPS III²². In this review, we will focus on recent developments for each therapy from their development in cellular and animal models, to the clinic.

3. The Road to the Clinic

3.1. Small molecules

3.1.2. Substrate reduction therapy

Substrate reduction therapy (SRT) aims to reduce the synthesis of GAGs, which cannot be degraded in MPS III. Unlike for other LSDs such as Gaucher type I, SRT is not approved for the treatment of any of the MPS disorders but several approaches to reducing GAG synthesis are under investigation²³. One approach is to silence the expression of genes which code for proteins involved in GAG synthesis using siRNA and shRNA as has been shown in MPS IIIA fibroblasts^{24,25} and MPS IIIC²⁶. An alternative approach to reducing the synthesis of GAG is the inhibition of follicle-stimulating hormone

(FSH) or epidermal growth factor (EGF), which have been previously shown to maximise the synthesis of some GAGs^{27,28}. Using a technique called gene expression-targeted isoflavone therapy (GET IT), Jakóbkiewicz-Banecka et al. showed that treatment of MPS IIIA and MPS IIIB patient-derived fibroblasts with the natural tyrosine kinase inhibitor isoflavone 4', 5, 7-trihydroxyisoflavone (genistein) inhibited GAG synthesis and prevented lysosomal accumulation. This effect was eliminated in the presence of excess EGF and partially restored following an increased concentration of genistein²⁹. In 2006, Malinowska et al. showed that continuous administration of genistein, given to MPSIIIB mice at a high dose for 9 months, significantly reduced lysosomal storage, heparan sulphate substrate and neuroinflammation in the cerebral cortex and hippocampus, resulting in correction of the behavioural defects observed, as well as improved synaptic vesicle protein expression and secondary storage in the cerebral cortex³⁰. As a small molecule, genistein can cross the blood-brain barrier and reach the brain, making it an attractive candidate for treating MPS IIIB.

An open-label clinical study including 19 MPS III patients (aged 2.8 – 19 years) was launched in 2014 to assess the safety and effectiveness of low dose genistein (5mg/kg/day) for one year and published by Delgado et al. Although no serious adverse effects were observed, the study showed that there was no improvement in the disability scale, as determined using the Questionnaire on Development and Behavior. Most patients had an increased disability score, or it remained the same despite a reduction in

urinary GAG levels³¹. As described above, mice treated with a high dose of genistein showed a significant reduction of HS accumulation and neuroinflammation in the brain and displayed an improvement in behaviour³⁰. Following these promising results, a phase III clinical trial was launched in 2013 using a higher-dose of genistein (160 mg/kg/day) or a placebo for a year, followed by a year of open-label genistein was completed recently (July 2018). In this double-blinded, randomized placebo-controlled study, high-dose (150 mg/kg/day) genistein was orally administered to nineteen MPS III patients (age 1.25 – 18.5, mean average age of 8) for a year (EudraCT number 2013–001479-18)³². Safety labs, GAG levels, clinical status and history of adverse events were obtained every 3 months, a physical examination was performed every 12 months, a 9 point disability scale (FPSS) was recorded after each visit, and an annual neurocognitive test was carried out where possible^{31,32}. After 12 months of treatment, no serious adverse events linked to high-dose genistein were identified and CSF HS was moderately reduced. However, the reduction in CSF was not substantial and scores on neuropsychological tests got worse or remained the same indicating no attenuation of cognitive decline³¹.

3.1.3. Other small molecules

Another way in which small molecules can be utilized to treat MPS III is by adopting strategies, which target the splicing process. 20% of MPS IIIC linked mutations reside within splice sites and affect mRNA processing, making this form of MPS an eligible candidate for this particular therapy

³³. In order to rescue the normal splicing process, Matos et al. used modified U1 snRNAs, which recognise mutations in donor splice sites in MPS IIIC patients' fibroblasts³³. Another approach, which targets mutations in an acceptor site and results in misfolded acetyl-CoA:α-glucosaminide acetyltransferase, tested a competitive inhibitor of the HGSNAT protein, glucosamine as a pharmacological chaperone to correct misfolded protein and restore normal trafficking to the lysosome³³. While partial correction of acetyl-CoA:α-glucosaminide acetyltransferase activity was achieved, the obstacle to fulfilling the full therapeutic potential of such a strategy is the efficient delivery of RNA molecules to the brain.

3.2. Enzyme Replacement Therapy

3.2.1. *In Vitro* studies

The blood-brain barrier has in the past been a major obstacle to effective intravenous treatment using ERT. This has been further confounded in MPS IIIB by inadequate mannose-6-phosphorylation (M6P) of human α-N-acetylglucosaminidase (rhNAGLU) recombinant enzyme, resulting in poor uptake. To address this problem, Kan et al. tested the use of a modified human recombinant NAGLU enzyme by fusing the human NAGLU fragment (rhNAGLU) to a fragment of insulin-like growth factor 2 (IGF-II) that would allow trafficking into the lysosome via IGFII binding site on the Mannose 6-phosphate/IGFII receptor. They successfully demonstrated that the fusion protein was able to gain entry to MPS IIIB cells via IGF-II binding to the mannose 6-

phosphate/IGFII receptor. The enzyme remained functional and reduced the amount of Heparan Sulphate (HS) in MPS IIIB fibroblasts to the same level as found in control cells^{34,35}.

This insulin-like growth factor II (IGFII)-tagged NAGLU molecule has been modified to create Tralesinidase alpha (rhNAGLU-IGFII; BMN 250), which will be further discussed in **section 3.2.3**. Yogalingam *et al.* (2019) used this fusion protein to distinguish two cellular uptake mechanisms by which BMN 250 is targeted to lysosomes in brain cells³⁶. Neurons, microglia and astrocytes are all critical cell types in MPS IIIB and therefore a better understanding of the cellular uptake mechanism(s) by which enzyme is delivered is important for developing efficient targeting mechanisms that will optimize ERT approaches. By systematically assessing the competitive cellular uptake of BMN 250 in human MPS IIIB patient fibroblasts and normal rodent-derived neurons, astrocytes and microglia, Yogalingam and colleagues found that BMN 250 is targeted to lysosomes in neurons, astrocytes and fibroblasts via MPR-mediated cellular uptake, whereas receptor-independent cellular uptake in microglia contributes to substantial lysosomal delivery of both BMN 250 and unfused rhNAGLU³⁶.

3.2.2. *In Vivo* studies

Intravenous delivery of ERT is already an available treatment for forms of MPS with less neurological involvement (MPS I³⁷, II³⁸, IVA³⁹, VI⁴⁰ and VII⁴¹). Adapting this classically systemic approach to gain entry to the CNS, has been the focus of recent studies for MPS IIIA and MPS IIIB. One way to

achieve this is by hijacking proteins, which have no trouble crossing the blood-brain barrier. One approach is to use a molecular Trojan horse, as has been achieved by fusing recombinant protein N-Sulfoglucosamine sulfohydrolase (SGSH) with a monoclonal antibody against the human insulin receptor (HIR Mab) (HIR Mab-SGSH) that can be taken up by MPS IIIA fibroblasts and trafficked to the lysosome resulting in reduced GAG levels (72 – 83%)⁴². Following intravenous administration of this fusion protein in Rhesus monkey, HIR Mab-SGSH 0.81% of the injected dose was detected in the brain⁴³. Similarly, Boado *et al.* have also created a fusion protein for MPS IIIB by fusing rhNAGLU to HIRMAb (HIRMAb-LL-NAGLU)⁴⁴. As shown for HIR Mab-SGSH in MPS IIIA, MPS IIIB fibroblasts displayed efficient cellular uptake of this fusion protein, trafficking to the lysosome, and a 74% reduction in the incorporation of sulfate into intracellular GAGs⁴⁴. When HIRMAb-LL-NAGLU was intravenously injected into a Rhesus monkey, 1% was detected in the brain⁴⁵.

A new promising *in vivo* study has recently been published, exploring the use of an IgG against mouse Transferrin Trojan Horse-Sulfamidase Fusion Protein in MPS IIIA mice. Large recombinant SGSH cannot cross the BBB, limiting intravenous administration a therapeutic option. To overcome this issue, Boado *et al.* have created a fusion protein consisting of an SGSH and IgG, where the IgG domain is a chimeric monoclonal antibody (MAb) against mouse transferrin receptor (TfR)⁴⁶. By acting as a Trojan horse, IgG can deliver SHSG to the CNS. The resulting fusion protein (cTfRMAb-SGSH)

successfully bound the mouse TfR with high affinity and maintained the same SGSH enzyme activity to the human recombinant SGSH. After 6 weeks of treatment, 3 times a week via intraperitoneal injection starting at 2 weeks of age, significant biochemical and behavioural improvements were observed. SGSH levels were elevated (36-fold compared to 30-fold in mice treated with just recombinant SGSH) and resulted in an 85% reduction in brain and liver HS (compared to 70% in mice treated with recombinant SGSH alone). Encouragingly, the reduction in brain HS was associated with a 28% increase in latency on the rotarod test of motor activity suggesting that an IgG-SGSH fusion protein engineered to penetrate the BBB via receptor-mediated transport, may be effective in treating MPS IIIA.

Another strategy that has been adopted for MPS IIIA and MPS IIIB involves direct administration of recombinant enzyme to the CNS. This was done in MPS IIIB by using recombinant human N-acetyl- α -glucosaminidase (rhNAGLU) fused with a fragment of the insulin-like growth factor II (IGF-II) as described in **section 3.2.1** and infusing it into the brain via intracerebroventricular injection ^{34,35}. Following intracerebroventricular administration to MPS IIIB mice, rhNAGLU-IGFII was taken up primarily by neurons and HS levels in the CSF were significantly reduced.

Similarly, when recombinant heparin N-sulfatase was infused to CSF of MPS IIIA dogs via the cisterna magna, HS levels were decreased in the CSF and cerebral cortex, but biomarkers linked to disease were only

normalized following high dose of enzyme, which as discussed previously, may have clinical implications in terms of such a treatment mounting an immune response ⁴⁷. Beard et al, have shown that the route of administration influences the success of ERT, as shown by infusing heparin N-sulfatase via Direct administration of recombinant enzyme to the brain can also be achieved via intrathecal lumbar, cisternal and ventricular administration. Lumbar infusion resulted in poor enzyme delivery and no significant reduction in GAG level, while infusion via the ventricular route proved more efficacious in decreasing GAG levels and dampening microglial activation ⁴⁸. Building on the relative success of this approach, the same group implanted an intraventricular cannula connected to a subcutaneous mini osmotic pump, allowing for a continuous low-dose infusion of recombinant human heparin N-sulfatase into the brain via the CSF. However, the therapeutic effects of this approach were not initially overwhelming, with only partial reduction of HS and GAG, and only moderate reductions in microglial activation but not astrogliosis ⁴⁹. By subsequently tweaking this method to improve implantation of the pumps in MPS IIIA, HS levels were normalized and GAG storage decreased significantly ⁵⁰.

Combinatorial approaches offer another way to improve the therapeutic potential of individual strategies. By combining the creation of fusion proteins with direct administration of enzyme directly into the brain, it is possible to further improve the impact on pathological biomarkers of either one therapy alone. This was demonstrated recently by Aoyagi-Scharber et al. in MPSIIIB mice. Human α -N-

acetylglucosaminidase fused with insulin-like growth factor II (described in **section 3.3.1**) and administered intracerebroventricularly resulted in widespread distribution of the fusion protein within the CNS and was accompanied by normalization of HS levels and significant reduction of secondary storage⁵¹. Although this combinatorial approach has yielded promising results, it is important to note the translational limitations surrounding frequent and direct administration of fusion proteins to patients.

3.2.3 Phase I/II Clinical Trials

Given the success of ERT in cellular and animal models, Shire Human Genetic Therapies (Shire HGT) developed an enzyme replacement therapy (ERT) recombinant human heparan-N-sulfatase (rhHNS) for patients with MPS IIIA. The open-label, phase I/II dose-escalation clinical study was carried out in twelve MPS IIIA patients to assess the safety of monthly intrathecal delivery of recombinant human heparin-N-sulfatase (rhHNS) using a surgically implanted intrathecal drug delivery device (IDDD) for the duration of 6 months (NCT01155778). In terms of safety, mild-to-moderate adverse effects were reported in all twelve patients, but none appeared to be related to the recombinant enzyme directly. However, despite a reduction in HS levels in the CSF, four of twelve patients showed a decline in the developmental quotient, six were stable and no dose group showed a clearly different response pattern. Overall, rhHNS administration via IDDD was generally safe and well-tolerated but required further investigation to determine efficacy⁵². Recently, an update on this study was

published by Wijburg *et al.* This phase I/II trial included twenty-one patients on a regimen of intrathecal rhHNS every two weeks, every 4 weeks or no treatment. Encouragingly, a clinical response to intrathecal rhHNS was observed in three of the treated patients. HS and GAG levels in the CSF were reduced in all treated patients. However, although treatment-emergent negative effects to intrathecal rhHNS were largely mild, no clear differences were detected between treated patients (age 17.8 – 47.8 months) and untreated controls (age 12.6 – 45.0 months) in terms of efficacy. Again, early intrathecal delivery of rhHNS is safe and effective at reducing HS and GAG levels in treated patients but the treatment has no neurocognitive effects (NCT02060526)⁵³.

The final results from a phase I/II, open-label, clinical study of intravenous recombinant human N-acetyl- α -D-glucosaminidase in children with mucopolysaccharidosis IIIB has recently been published. The study, sponsored by Alexicon Pharmaceuticals, included 11 participants age between 1 – 10 years of age and set out to evaluate the safety, tolerability, pharmacokinetics, and efficacy of intravenous administration of a SBC-103, a recombinant human NAGLU enzyme capable of crossing the blood-brain barrier⁵⁴ (NCT02324049). In this three-part study, participants were sequentially divided into three dose-escalating groups and received intravenous injections every two weeks for 24 weeks, after which patients received no treatment for a month (Part I). Patients then received a higher dose every two weeks (Part II) starting at 28 weeks, and a final dose escalation of SBC-103 every two weeks for two years in total⁵⁵. Despite the initial results in NAGLU-deficient

mice ⁵⁴, SBC-103 (rhNAGLU) was well-tolerated by MPS IIIB patients, and resulted in the reduction of HS in the CSF but had no effect on preventing brain atrophy or preventing neurocognitive decline for at any dose. Interestingly, SBC-103 was not detected in the CSF suggesting that it may not have reached the CNS⁵⁵.

3.3. Gene therapy

3.3.1. Adeno-associated virus (AAV)

To overcome the need for periodic administration of ERT or therapies based on small molecules like SRT, one-shot gene therapies are being developed for MPS III in order to provide constant production of the deficient enzyme. One of the main advantages of gene therapy is that only a proportion of cells in an organ, in this case, the brain, need to be corrected, as these corrected cells can produce a sufficient amount of the active enzyme to cross-correct neighbouring cells ⁵⁶⁻⁵⁹. Arguably, gene therapy approaches currently stand out as the most promising therapeutic approach for treating MPS III, with the publication of various studies demonstrating the potential use of gene therapy to treat not just MPS III, but numerous other monogenic neurological diseases ⁶⁰. However, despite this promise, there are still hurdles to therapeutic efficacy with this type of therapy, relating mainly to vector type and route of administration. For example, are there differences in transduction efficiency between adeno-associated virus (AAV) and lentivirus? Can intravenously administered viruses infect neurons in the brain and is the direct administration of vector to the brain is safe?

Many AAV-based strategies involve the delivery of genes to the CNS. To achieve widespread distribution, initial experiments adopting AAV used multiple direct injections to the brain parenchyma. As the cell and tissue tropism of different AAV serotypes (AAV1 – AAV13) became better understood ⁶¹, subsequent experiments began to take advantage of the ability of certain AAV serotypes to gain access to the CNS. For example, AAV9 can cross the BBB after intravenous administration, resulting in widespread transduction of the CNS and the peripheral organs through a non-invasive procedure. Alternatively, various other AAV vectors have been administered directly to the CSF, hijacking the ventricular system to achieve global CNS gene transfer, as well as delivery to the peripheral nervous system and liver. In this section, we will focus on various studies, which together demonstrate the efficacy of different vectors and routes of administration for the treatment of MPS III.

Most gene therapy studies on MPS III use the vector adeno-associated virus (AAV), which achieve high transduction *in vivo* and have proven safe in clinical studies ^{62,63}. Furthermore, preclinical and clinical data provide evidence for long-term AAV-mediated gene expression in the brain, without producing any significant adverse effect ^{56,58,64-69}. However, in terms of efficacy, the selection of AAV virus serotype is a crucial consideration. Gilkes *et al.* have used AAV5, AAV8, AAV9 and AAVrh10 to deliver NAGLU to MPS IIIB mice via direct administration of the virus to the CNS. Although they found that AAV8 showed the greatest efficacy in terms of bio-distribution and transduction of NAGLU ⁷⁰, other studies

using AAV9-mediated NAGLU gene transfer via the CSF or systemic delivery, have also proved efficacious. A study by Ribera *et al.* administered AAV9 vectors carrying NAGLU to the CSF of MPS IIIB mice at 2 months of age when the disease has already become established and observed a restoration of gene expression and enzymatic activity in the CNS, which resulted in normalization of GAGs and lysosomal physiology and reduced neuroinflammation. The systemic hallmarks of the disease were also corrected, behavioural deficits improved and lifespan was extended⁵⁸. More recently, the metabolomics profiles of MPS IIIB mice was specifically measured in MPS IIIB mice to assess the impact of systemic gene delivery⁷¹. Following intravenous administration of rAAV9-hNAGLU, near-complete correction of systemic metabolomic impairments was observed⁷¹. Ahead of a clinical gene therapy treatment, an AAV9-mediated vector carrying human NAGLU (rAAV9-CMV-hNAGLU) was tested on cynomolgus monkeys via intravenous injection⁷². Over the course of 6 months, no adverse effects were apparent and the treatment resulted in long-lasting global CNS and somatic transduction, with relatively high NAGLU activity compared to wildtype levels, both in the brain and body⁷². This study demonstrated an effective and safe profile for systemic rAAV9-hNAGLU vector delivery in nonhuman primates, providing evidence for its clinical potential in humans. Safety of intravenous administration of AAV9-mediated NAGLU gene transfer via the CMV promoter (rAAV9-CMV-hNAGLU) has also been tested. Meadows *et al.* performed an IND-enabling good laboratory practice (GLP) toxicology study in healthy

and MPS IIIB mice. rNAGLU expression was rapid and persistent in the majority of CNS without any adverse clinical signs of toxicology during the 6-month study, but a dosing range for safe and effective rAAV9-CMV-hNAGLU systemic gene delivery in MPS IIIB was also identified⁷³.

Different AAV serotypes have also been tested in other forms of MPS III. AAVrh10 has been used to deliver SGSH in MPS IIIA mice via intraparenchymal administration⁷⁴. AAVrh10-derived SGSH enzyme improved the breakdown of heparan sulfate and reduced microglial activation. With time, GM3 ganglioside accumulation was ameliorated and the formation of ubiquitin-positive lesions near the injection site or in regions connected to the injection site was prevented. However, these positive changes were restricted to the site of injection, and no such changes were observed in regions of the brain distant from or lacking connections with, the administration site. Therefore, to obtain adequate therapeutic efficacy, it may be necessary to administer the gene vector to multiple intraparenchymal regions in order to ensure widespread distribution of enzyme and correction of disease pathology, increasing the likelihood of infection⁷⁴. Recently, AAV9 was used to test gene therapy in the MPS IIID mouse model for the first time⁵⁹. Treatment of the GNS-deficient animals with GNS-expressing AAV9-derived vector delivered to the cerebrospinal fluid normalized GAG storage, improved lysosomal functionality in the CNS and somatic tissues, reduced neuroinflammation, restored normal behaviour and extended the lifespan of treated mice relative to untreated MPS IIID mice⁵⁹.

3.3.2. Autoantibodies to Adeno-associated virus

One commonly cited drawback of the use of AAV-mediated gene therapy is that individuals with pre-existing host humoral and cellular immunity to AAV capsids may be subject to limited target tissue transduction and long-term expression of the genes they carry, making them less likely to benefit from AAV-mediated gene transfer ⁷⁵. Even low levels of neutralizing antibodies against AAV capsids can result in impaired transduction of the incorporated gene following intravenous delivery ^{76,77}. However, Murrey et al. showed that low levels of preexisting anti-AAV9 antibodies did not affect vector transduction of rAAV9-CMV-hNAGLU in cynomolgus monkeys. Even at high levels, preexisting anti-AAV9 Abs led to reduced transduction in the liver and other somatic tissues but did not diminish transgene expression in the brain ⁷². Similarly, Ribera et al. showed in their study that enzymatic activity in the CSF of dogs after administration of canine NAGLU-coding vectors to animals that were either naïve or had pre-existing immunity against AAV9, displayed similar levels of enzyme activity, suggesting that CNS efficacy would not be impaired in patients that are seropositive for AAV9 ⁵⁸. These studies demonstrate that at least for AAV9, an effective and safe profile for systemic vector delivery in nonhuman primates can be achieved.

3.3.3. Lentiviral/Adeno-associated virus combinatorial approach

Like AAV-mediated gene therapy, lentivirus has proved effective in treating MPS III.

Lentiviral vectors carrying genes coding for murine heparin N-sulfatase and sulfatase modifying factor-1 have been tested in MPS IIIA. After administration via the cerebral lateral ventricles, enzyme activity was found to be between 0.5- and 4-fold greater than in normal mouse brain and ganglioside and lysosomal β -hexosaminidase levels, both of which are characteristically elevated in MPS IIIA, were significantly reduced, or were normalised ⁷⁸. Furthermore, combining different vector types via alternative routes can prove more effective at achieving disease correction than either one alone. In one study, AAV2/5-mediated and lentivirus-mediated NAGLU expression was more efficient than either one therapy alone in treating MPS IIIB ⁷⁹. MPS IIIB neonatal mice were treated with intracranial AAV2/5-NAGLU, intravenous lentiviral-NAGLU or both. All treatment groups resulted in significant biochemical and histological improvements compared with untreated MPS IIIB animals, but the animals treated with both AAV2/5 and lentivirus lived significantly longer (612 days) than animals treated with just AAV2/5-mediated gene therapy (463) or lentiviral gene therapy (358) suggesting that although MPS III disease is primarily neurological, targeting both the systemic and central nervous system early in life appears to be the most efficacious approach for treating MPS IIIB ⁷⁹.

3.3.4. Phase I/II Clinical trials

A number of gene therapy clinical trials have been sponsored to test the safety and efficacy of gene therapy for the treatment of MPS III. In this review, we will focus on studies which have available data but a complete overview is described in detail by Marco et al. ⁶⁰. These

studies include phase I/II trials for MPS IIIA, MPS IIIB and MPS IIIC. To date, only one clinical trial has been completed, a phase I/II trial testing intracerebral administration of AAV10 carrying the human SGSH and SUMF1 cDNAs (SAF-301; rh.10-SGSH-IRES-SUMF1) for the treatment of MPS IIIA (NCT01474343 and NCT02053064). In this study, Lysogene recruited four children (three aged 5.5 – 6 years old, and one aged 2 years 8 months) to test the tolerance and safety of SAF-301, and assess disease biomarkers in blood, urine and CSF and brain function during one year of follow up. The results were published by Tardieu *et al.*⁸⁰ and reported that the therapy was safe and well-tolerated and improved brain atrophy and behaviour. All children showed a decline in cognitive ability and three patients presented with brain atrophy. After 8 weeks of treatment, MRI showed that brain atrophy has stabilized in two patients but increased in the other two, and there was a moderate improvement in behaviour, attention and sleep in three of the patients⁸⁰. An open-label long term study was initiated five years after treatment to follow up on patients with MPS IIIA who had previously been treated with SAF-301 which ended in 2017, but no results were available. The aim was to collect additional safety and tolerability data on the treatment, and further collect data to assess the effects of SAF-301 on neurological and psychological status and biomarkers (NCT02053064).

A number of clinical studies are still underway, Esteve has developed EGT-101, a compound consisting of AAV9 containing hSGSH (AAV9-hSGSH). In this phase I/II clinical trial, EGT-101 has been administered via intra-CSF administration to MPS IIIA

patients (2015-000359-26). An uncontrolled phase I/II clinical trial sponsored by UniQure Biopharma is also currently investigating the intraparenchymal administration of a recombinant AAV2/5 vector encoding human NAGLU AAV5-hNAGLU in four MPS IIIB patients (NCT03300453), the results of which have recently been published⁸¹. 30 months after injection, the treatment appeared to be safe and well-tolerated with sustained NAGLU production (15-20% of that in unaffected children) in the CSF. Compared with the natural history of MPS III syndromes, neurocognitive progression was improved in all patients, with the youngest patient having function comparable to that in healthy children.

Abeona Therapeutics is currently recruiting for two phase I/II clinical trials for both MPS IIIA and MPS IIIB. To treat MPS IIIA, a self-complementary AAV9 vector carrying the human SGSH gene under the control of a U1a promoter (scAAV9.U1a.hSGSH) called ABO-102 was delivered intravenously to participants two years of age or older in an open-label, dose-escalation phase I/II clinical trial (NCT02716246). The estimated number of participants for this study is 22 and the primary aim is to assess safety and neurocognitive function (developmental score) after 24 months and secondarily, to assess SGSH activity, liver and spleen volume, cognitive ability and urinary GAG levels. So far, no adverse events relating to scAAV9.U1a.hSGSH have been reported. Although efficacy data is yet to be published, some preliminary data is available, showing a dose-dependent and sustained reduction in CSF HS all three cohorts after 30 days. Following dosing at 14-26 months of age,

participants showed normal development 12 – 18 months post-treatment. To treat MPS IIIB, Abeona Therapeutics have also sponsored a new phase I/II trial using one-time intravenous administration of AAV9 carrying the human NAGLU gene under the control of a CMV enhancer/promoter (rAAV9.CMV.hNAGLU) called ABO-101 (NCT03315182). This two-year open-labelled, dose-escalation clinical trial will include an estimated 9 MPS IIIB patients aged between 6 – 2 years or older with a minimum Developmental Quotient of 60 or above. Two doses (2×10^{13} vg/kg and 5×10^{13} vg/kg) are being tested across two cohorts to primarily assess safety and neurodevelopment, and other secondary endpoints including neurocognitive and behavior evaluations, quality of life, enzyme activity in cerebrospinal fluid (CSF) and plasma, biomarkers in CSF, plasma and urine, and brain and liver volume. As described above, the preliminary data from the MPS IIIA study are encouraging, but no data is yet available for MPS IIIB⁶⁰ (www.clinicaltrials.org). While gene therapy has, and continues to show tremendous therapeutic promise for the treatment of MPS III disease in animal models, leading to a number of clinical trials, these trials have revealed that the ideal method of delivery of viral vectors is yet to be elucidated.

3.4. Cellular Therapies

3.4.1. Hematopoietic Stem Cells

Hematopoietic Stem Cells treatment (HSCTs) can be obtained from the bone marrow or peripheral blood of a healthy donor and transplanted into a patient. To avoid the

rejection of donor cells by the patient's immune system, they must first be immunosuppressed according to a conditioning regimen⁸². Healthy, matched enzyme-secreting donor cells can then be transplanted into the patient, providing a permanent and continuous supply of protein. For a more detailed description of the history and application of HSCs in MPS, Taylor et al. have written a thorough review⁸³. HSCTs is already an effective therapy for a number of inborn errors of metabolism, a good example of which is in the treatment of Hurler's syndrome if administered early. However, what has been learnt from the clinical application of HSCTs for the treatment of Hurler Syndrome, has not translated well to the treatment of MPS III⁸⁴. Unfortunately, HSCT have not proved successful in preventing the progression of neurological disease in MPS III patients.

Furthermore, even when HSCT is administered to MPS IIIB patients before the onset of neurological symptoms, studies have shown that neurocognitive decline still ensues^{85,86}. A clinical study was performed in 62 MPS patients, only 2 of which had been diagnosed with MPS III, and found that although the treatment was safe and effective overall in MPS, it is difficult to conclude the efficacy to MPS III specifically⁸⁷.

3.4.2. Umbilical Cord Mononuclear Cells

A large study on unrelated donor umbilical cord blood transplantation for inherited metabolic disorders in 159 patients, showed more promise than previously described in **section 3.4.1**. Of the 19 MPS III patients enrolled in the study, 12 survived and 9

showed disease stabilization with lesser neurological symptoms. Overall, children who received HSCT appeared to have fewer behavioral problems and better sleeping patterns as compared with children who did not receive transplants. One MPS IIIB patients received HSCT just before their second birthday and appeared to respond best to the treatment. At age 15, the patient had normal blood levels of O sulphated HS and N-sulfated HS and disease symptoms appeared to be better than MPS IIIB patients who had not received treatment⁸⁸. However, in another study, umbilical cord blood-derived hematopoietic stem cells (UCBT) were transplanted into two MPS III patients (MPS IIIA and MPS IIIB) before the onset of neurological symptoms and monitored for 5-years. Despite uncomplicated transplantation, with full engraftment of donor cells, both patients showed progressive neurological deterioration, regression of cognitive skills, and behavioural disturbances, which was comparable to untreated patients with the same mutations⁸⁹. In addition, the HS concentration in CSF in the MPS IIIB patient was just as high as in untreated MPS IIIB patients. Given the outcome of this clinical study, it can be concluded that like BMT, early UCBT does not prevent neurological deterioration in MPS III.

Although attempts at using HSC and UCBT transplantations in MPS III have yielded disappointing results in recent years, leading to the assumption that these approaches have little potential for being effective treatments^{3,12,13}, animal studies have continued to raise hope. In the mouse models of MPS IIIB, monthly intravenous administration of human umbilical cord mononuclear cells over a

period of six months proved effective in reducing ganglioside accumulation, microglial activation, corrected anxiety-like behaviour and restored hippocampal cytoarchitecture⁹⁰.

3.4.4. Ex-vivo gene modification

In vivo gene therapy strategies involve the administration of viral vector particles directly to patients to provide affected cells with normal complementary DNA, *ex vivo* gene therapy approaches are based on the *ex vivo* transduction of patient cells that are subsequently infused back, potentially circumventing an immune response for foreign cells⁹¹. For a more extensive comparison between *in vivo* and *ex vivo* approaches to date, Fraldi et al. have recently published a review. In brief, gene therapy approaches have been developed for MPS IIIA and MPS IIIB using autologous transplantation of HSCs genetically modified using lentiviral vectors to express SGSH or NAGLU⁹²⁻⁹⁵. After transplantation, gene-corrected cells proliferate and travel to the brain where they cross the BBB to become resident cells of the CNS. Here, they secrete the deficient protein, and subsequently cross-correct other endogenous cells. These studies have shown normalization of HS, secondary storage and neuroinflammation as well as improvements in behavioural read-outs.

One way to overcome the blood-brain barrier is to target therapy directly to the CNS via parenchymal injection or via the cerebrospinal fluid. In recent years, Clarke et al. have shown that NSCs derived from reprogrammed MPS IIIB mouse embryonic fibroblasts to create iPSCs and corrected using lentiviral-

mediated human NAGLU overexpression, alleviated neuropathology⁹⁶. It is important to underline the fact that a modified NAGLU enzyme was not necessary, as described in **section 3.2.1** above. Furthermore, very little enzyme was needed to obtain correction. These findings suggest that cell therapies represent an important line of investigation, despite current dogma. The use of neural progenitor cells to provide the missing enzyme also has regenerative potential since they can differentiate *in vivo* into neurons and astrocytes⁹⁷⁻¹⁰¹. This is currently an underdeveloped field, given the fact that attempts to promote regeneration in the spinal cord injuries have been challenging. Nevertheless, several reports from studies on neurodegenerative animal models show that neural progenitor cells can differentiate into functional neurons, which are capable of restoring neuronal networks to a degree that impacts neurocognition behaviour^{97,99,101}. For Parkinson disease, the use of neural progenitor cells is further along in the developmental process and several clinical trials are underway addressing safety and efficacy to treat Parkinson disease (NCT03128450, NCT03128450, NCT03309514, NCT03815071, NCT02452723, NCT02452723). Finally it important to underline that despite the obstacles posed by stem cell therapies compared to others, this line of research should be further pursued as it offers a unique opportunity to address neuronal loss, which other therapies do not.

CONCLUSION

The genetic cause of MPS III and the biochemistry of their gene products are well

known and methods for genetic and biochemical diagnosis have been established. Therapeutic approaches have been developed, which target key aspects of the disease from its root cause to its downstream effects. With the same knowledge, HSCTs and ERT have been developed for the treatment of other forms of MPS and now both have been approved for MPS I, II, IVA, VI and VII. However, for MPS III, the road to therapy has been and continues to be a challenge given its neurological nature. Fantastic progress has been made in adapting therapies for other forms of MPS, offering up important lessons. Cellular and animal models have paved the way for several clinical trials. Among them the most advanced in development are ERT, involving either direct administration to the brain or the use of BBB-compliant fusion protein, and gene therapy using vectors administered either direct to the brain or via the bloodstream like ERT. Both approaches have presented new challenges at clinical trials, but new and modified approaches are currently under investigation. Although these approaches stand out as the most advanced, it is important to recognise their limitations now and in the future, and remain open to overcoming the barriers to other forms of therapy such as stem cell therapies. While stem cell therapies have proved disappointing to date, research into stem cell treatment for other neurodegenerative disease continue to show promise and in the process reveal areas for improvement that may be applicable to MPS III. Another important consideration, when asking how close we are to a therapy for MPS III is whether one therapy alone will ever be enough. Even if ERT, substrate reduction therapy, gene therapy or stem cell

therapies are optimised, it may be necessary to combine different approaches in order to further improve pathological outcome measures and behavioural phenotypes, but most importantly, extend the life span of patients.

A frustration in translational research is why promising results in non-human models often lead to disappointing results in clinical trials. It is worth pointing out that MPS III is a rare disorder making it difficult to normalise studies for treatment groups while maintaining a high enough number of participants to generate meaningful data. It is also important to point out that there is a disproportionate number of studies conducted on MPS IIIA and MPS IIIB. This means that

some therapies, such as ERT and gene therapy, may become available for these subtypes more quickly than others. Nonetheless, despite its low prevalence, MPS III is a severely debilitating disease affecting not only the patient, parents and caregivers but society, justifying further attention and research. While researchers continue to develop therapies for MPS III, multidisciplinary teams who consider the age, clinical stage, severity and socioeconomic status of patients are essential for the proper management of those suffering with MPSIII. Non-profit organization have a pivotal role in promoting initial studies for therapy development, but this effort should be further supported by governments and pharmaceutical companies.

Reference

1. Andrade F, Aldamiz-Echevarria L, Llarena M, Couce ML. Sanfilippo syndrome: Overall review. *Pediatr. Int.* Jun 2015;57(3):331-338.
2. Fedele AO. Sanfilippo syndrome: causes, consequences, and treatments. *The application of clinical genetics.* 2015;8:269-281.
3. Jakobkiewicz-Banecka J, Gabig-Ciminska M, Kloska A, et al. Glycosaminoglycans and mucopolysaccharidosis type III. *Frontiers in bioscience (Landmark edition).* Jun 1 2016;21:1393-1409.
4. Kresse H, Neufeld EF. The Sanfilippo A corrective factor. Purification and mode of action. *J. Biol. Chem.* Apr 10 1972;247(7):2164-2170.
5. von Figura K. Human alpha-n-acetylglucosaminidase. 2. Activity towards natural substrates and multiple recognition forms. *Eur. J. Biochem.* Nov 1 1977;80(2):535-542.
6. Klein U, Kresse H, von Figura K. Sanfilippo syndrome type C: deficiency of acetyl-CoA:alpha-glucosaminide N-acetyltransferase in skin fibroblasts. *Proc. Natl. Acad. Sci. U. S. A.* Oct 1978;75(10):5185-5189.
7. Kowalewski B, Lamanna WC, Lawrence R, et al. Arylsulfatase G inactivation causes loss of heparan sulfate 3-O-sulfatase activity and mucopolysaccharidosis in mice. *Proc. Natl. Acad. Sci. U. S. A.* Jun 26 2012;109(26):10310-10315.
8. Valstar MJ, Ruijter GJ, van Diggelen OP, Poorthuis BJ, Wijburg FA. Sanfilippo syndrome: a mini-review. *J. Inherit. Metab. Dis.* Apr 2008;31(2):240-252.
9. Cross EM, Hare DJ. Behavioural phenotypes of the mucopolysaccharide disorders: a systematic literature review of cognitive, motor, social, linguistic and behavioural presentation in the MPS disorders. *J. Inherit. Metab. Dis.* Mar 2013;36(2):189-200.
10. Lyon GK, E.H. Pastores, G.M. *Neurology of Hereditary Metabolic Diseases of Children.* 3 ed. New York: Mc Graw Hill; 2006.
11. Wraith JE. Mucopolysaccharidoses and Oligosaccharidoses. In: Fernandes J, Saudubray J-M, van den Berghe G, Walter JH, eds. *Inborn Metabolic Diseases: Diagnosis and Treatment.* Berlin, Heidelberg: Springer Berlin Heidelberg; 2006:495-507.
12. Giugliani R, Federhen A, Vairo F, et al. Emerging drugs for the treatment of mucopolysaccharidoses. *Expert opinion on emerging drugs.* 2016;21(1):9-26.

- 13.** Gaffke L, Pierzynowska K, Piotrowska E, Wegrzyn G. How close are we to therapies for Sanfilippo disease? *Metab. Brain Dis.* Feb 2018;33(1):1-10.
- 14.** Lancaster MA, Knoblich JA. Generation of cerebral organoids from human pluripotent stem cells. *Nat. Protoc.* Oct 2014;9(10):2329-2340.
- 15.** Allende ML, Cook EK, Larman BC, et al. Cerebral organoids derived from Sandhoff disease-induced pluripotent stem cells exhibit impaired neurodifferentiation. *J. Lipid Res.* Mar 2018;59(3):550-563.
- 16.** Camp JG, Badsha F, Florio M, et al. Human cerebral organoids recapitulate gene expression programs of fetal neocortex development. *Proc. Natl. Acad. Sci. U. S. A.* Dec 22 2015;112(51):15672-15677.
- 17.** Cederquist GY, Asciolla JJ, Tchieu J, et al. Specification of positional identity in forebrain organoids. *Nat. Biotechnol.* Apr 2019;37(4):436-444.
- 18.** Choi SH, Kim YH, Hebisch M, et al. A three-dimensional human neural cell culture model of Alzheimer's disease. *Nature.* Nov 13 2014;515(7526):274-278.
- 19.** Jo J, Xiao Y, Sun AX, et al. Midbrain-like Organoids from Human Pluripotent Stem Cells Contain Functional Dopaminergic and Neuromelanin-Producing Neurons. *Cell stem cell.* Aug 4 2016;19(2):248-257.
- 20.** Vallejo S, Fleischer A, Martin JM, Sanchez A, Palomino E, Bachiller D. Generation of two induced pluripotent stem cells lines from Mucopolysaccharydosis IIIA patient: IMEDEAi004-A and IMEDEAi004-B. *Stem cell research.* Oct 2018;32:110-114.
- 21.** Vallejo-Diez S, Fleischer A, Martin-Fernandez JM, Sanchez-Gilabert A, Bachiller D. Generation of two induced pluripotent stem cells lines from a Mucopolysaccharydosis IIIB (MPSIIIB) patient. *Stem cell research.* Dec 2018;33:180-184.
- 22.** Ghosh A, Shapiro E, Rust S, et al. Recommendations on clinical trial design for treatment of Mucopolysaccharidosis Type III. *Orphanet J. Rare Dis.* Jun 26 2017;12(1):117.
- 23.** Platt FM, Butters TD. Substrate Reduction Therapy. *Lysosomal Storage Disorders.* Boston, MA: Springer US; 2007:153-168.
- 24.** Dziezic D, Wegrzyn G, Jakobkiewicz-Banecka J. Impairment of glycosaminoglycan synthesis in mucopolysaccharidosis type IIIA cells by using siRNA: a potential therapeutic approach for Sanfilippo disease. *Eur. J. Hum. Genet.* Feb 2010;18(2):200-205.
- 25.** Kaidonis X, Liaw WC, Roberts AD, Ly M, Anson D, Byers S. Gene silencing of EXTL2 and EXTL3 as a substrate deprivation therapy for heparan sulphate storing mucopolysaccharidoses. *Eur. J. Hum. Genet.* Feb 2010;18(2):194-199.
- 26.** Canals I, Beneto N, Cozar M, Vilageliu L, Grinberg D. EXTL2 and EXTL3 inhibition with siRNAs as a promising

substrate reduction therapy for Sanfilippo C syndrome. *Sci. Rep.* Sep 8 2015;5:13654.

27. Pisano MM, Greene RM. Epidermal growth factor potentiates the induction of ornithine decarboxylase activity by prostaglandins in embryonic palate mesenchymal cells: effects on cell proliferation and glycosaminoglycan synthesis. *Dev. Biol.* Aug 1987;122(2):419-431.

28. Tirone E, D'Alessandris C, Hascall VC, Siracusa G, Salustri A. Hyaluronan synthesis by mouse cumulus cells is regulated by interactions between follicle-stimulating hormone (or epidermal growth factor) and a soluble oocyte factor (or transforming growth factor beta1). *J. Biol. Chem.* Feb 21 1997;272(8):4787-4794.

29. Jakobkiewicz-Banecka J, Piotrowska E, Narajczyk M, Baranska S, Wegrzyn G. Genistein-mediated inhibition of glycosaminoglycan synthesis, which corrects storage in cells of patients suffering from mucopolysaccharidoses, acts by influencing an epidermal growth factor-dependent pathway. *J. Biomed. Sci.* Mar 2 2009;16:26.

30. Malinowska M, Wilkinson FL, Langford-Smith KJ, et al. Genistein improves neuropathology and corrects behaviour in a mouse model of neurodegenerative metabolic disease. *PLoS One.* Dec 1 2010;5(12):e14192.

31. Delgadillo V, O'Callaghan Mdel M, Artuch R, Montero R, Pineda M. Genistein supplementation in patients affected by Sanfilippo disease. *J. Inherit. Metab. Dis.* Oct 2011;34(5):1039-1044.

32. Kim KH, Dodsworth C, Paras A, Burton BK. High dose genistein aglycone therapy is safe in patients with mucopolysaccharidoses involving the central nervous system. *Mol. Genet. Metab.* Aug 2013;109(4):382-385.

33. Matos L, Canals I, Dridi L, et al. Therapeutic strategies based on modified U1 snRNAs and chaperones for Sanfilippo C splicing mutations. *Orphanet J. Rare Dis.* Dec 10 2014;9:180.

34. Kan SH, Aoyagi-Scharber M, Le SQ, et al. Delivery of an enzyme-IGFII fusion protein to the mouse brain is therapeutic for mucopolysaccharidosis type IIIB. *Proc. Natl. Acad. Sci. U. S. A.* Oct 14 2014;111(41):14870-14875.

35. Kan SH, Troitskaya LA, Sinow CS, et al. Insulin-like growth factor II peptide fusion enables uptake and lysosomal delivery of alpha-N-acetylglucosaminidase to mucopolysaccharidosis type IIIB fibroblasts. *Biochem. J.* Mar 01 2014;458(2):281-289.

36. Yogalingam G, Luu AR, Prill H, et al. BMN 250, a fusion of lysosomal alpha-N-acetylglucosaminidase with IGF2, exhibits different patterns of cellular uptake into critical cell types of Sanfilippo syndrome B disease pathogenesis. *PLoS One.* 2019;14(1):e0207836.

37. Kakkis ED, Muenzer J, Tiller GE, et al. Enzyme-replacement therapy in mucopolysaccharidosis I. *N. Engl. J. Med.* Jan 18 2001;344(3):182-188.

- 38.** Muenzer J, Guzsavos-Calikoglu M, McCandless SE, Schuetz TJ, Kimura A. A phase I/II clinical trial of enzyme replacement therapy in mucopolysaccharidosis II (Hunter syndrome). *Mol. Genet. Metab.* Mar 2007;90(3):329-337.
- 39.** Hendriksz CJ, Burton B, Fleming TR, et al. Efficacy and safety of enzyme replacement therapy with BMN 110 (elosulfase alfa) for Morquio A syndrome (mucopolysaccharidosis IVA): a phase 3 randomised placebo-controlled study. *J. Inherit. Metab. Dis.* Nov 2014;37(6):979-990.
- 40.** Harmatz P, Giugliani R, Schwartz I, et al. Enzyme replacement therapy for mucopolysaccharidosis VI: a phase 3, randomized, double-blind, placebo-controlled, multinational study of recombinant human N-acetylgalactosamine 4-sulfatase (recombinant human arylsulfatase B or rhASB) and follow-on, open-label extension study. *J. Pediatr.* Apr 2006;148(4):533-539.
- 41.** Fox JE, Volpe L, Bullaro J, Kakkis ED, Sly WS. First human treatment with investigational rhGUS enzyme replacement therapy in an advanced stage MPS VII patient. *Mol. Genet. Metab.* Feb 2015;114(2):203-208.
- 42.** Boado RJ, Lu JZ, Hui EK, Pardridge WM. Insulin receptor antibody-sulfamidase fusion protein penetrates the primate blood-brain barrier and reduces glycosaminoglycans in Sanfilippo type A cells. *Mol. Pharm.* Aug 4 2014;11(8):2928-2934.
- 43.** Boado RJ, Ka-Wai Hui E, Zhiqiang Lu J, Pardridge WM. Insulin receptor antibody-iduronate 2-sulfatase fusion protein: pharmacokinetics, anti-drug antibody, and safety pharmacology in Rhesus monkeys. *Biotechnol. Bioeng.* Nov 2014;111(11):2317-2325.
- 44.** Boado RJ, Lu JZ, Hui EK, Lin H, Pardridge WM. Insulin Receptor Antibody-alpha-N-Acetylglucosaminidase Fusion Protein Penetrates the Primate Blood-Brain Barrier and Reduces Glycosaminoglycans in Sanfilippo Type B Fibroblasts. *Mol. Pharm.* Apr 4 2016;13(4):1385-1392.
- 45.** Boado RJ, Hui EK, Lu JZ, Pardridge WM. Very High Plasma Concentrations of a Monoclonal Antibody against the Human Insulin Receptor Are Produced by Subcutaneous Injection in the Rhesus Monkey. *Mol. Pharm.* Sep 6 2016;13(9):3241-3246.
- 46.** Boado RJ, Lu JZ, Hui EK, Pardridge WM. Reduction in Brain Heparan Sulfate with Systemic Administration of an IgG Trojan Horse-Sulfamidase Fusion Protein in the Mucopolysaccharidosis Type IIIA Mouse. *Mol. Pharm.* Feb 5 2018;15(2):602-608.
- 47.** King B, Marshall N, Beard H, et al. Evaluation of enzyme dose and dose-frequency in ameliorating substrate accumulation in MPS IIIA Huntaway dog brain. *J. Inherit. Metab. Dis.* Mar 2015;38(2):341-350.
- 48.** Beard H, Luck AJ, Hassiotis S, et al. Determination of the role of injection site on the efficacy of intra-CSF enzyme replacement

therapy in MPS IIIA mice. *Mol. Genet. Metab.* May 2015;115(1):33-40.

49. King B, Setford ML, Hassiotis S, et al. Low-dose, continual enzyme delivery ameliorates some aspects of established brain disease in a mouse model of a childhood-onset neurodegenerative disorder. *Exp. Neurol.* Apr 2016;278:11-21.

50. King B, Hassiotis S, Rozaklis T, et al. Low-dose, continuous enzyme replacement therapy ameliorates brain pathology in the neurodegenerative lysosomal disorder mucopolysaccharidosis type IIIA. *J. Neurochem.* May 2016;137(3):409-422.

51. Aoyagi-Scharber M, Crippen-Harmon D, Lawrence R, et al. Clearance of Heparan Sulfate and Attenuation of CNS Pathology by Intracerebroventricular BMN 250 in Sanfilippo Type B Mice. *Molecular therapy. Methods & clinical development.* Sep 15 2017;6:43-53.

52. Jones SA, Breen C, Heap F, et al. A phase 1/2 study of intrathecal heparan-N-sulfatase in patients with mucopolysaccharidosis IIIA. *Mol. Genet. Metab.* Jul 2016;118(3):198-205.

53. Wijburg FA, Whitley CB, Muenzer J, et al. Intrathecal heparan-N-sulfatase in patients with Sanfilippo syndrome type A: A phase IIB randomized trial. *Mol. Genet. Metab.* Feb 2019;126(2):121-130.

54. Rutkowski JV, Harbert K, Xu H, et al. Intravenous SBC-103, a recombinant human alpha-N-acetylglucosaminidase reduces CNS heparan sulfate content in a

mucopolysaccharidosis type IIIB mouse model. *Mol. Genet. Metab.* 2014/02/01/2014;111(2):S92.

55. Whitley CB, Vijay S, Yao B, et al. Final results of the phase 1/2, open-label clinical study of intravenous recombinant human N-acetyl-alpha-d-glucosaminidase (SBC-103) in children with mucopolysaccharidosis IIIB. *Mol. Genet. Metab.* Feb 2019;126(2):131-138.

56. Haurigot V, Marco S, Ribera A, et al. Whole body correction of mucopolysaccharidosis IIIA by intracerebrospinal fluid gene therapy. *J. Clin. Invest.* Aug 1 2013;123(8):3254-3271.

57. Hocquemiller M, Giersch L, Audrain M, Parker S, Cartier N. Adeno-Associated Virus-Based Gene Therapy for CNS Diseases. *Hum. Gene Ther.* Jul 2016;27(7):478-496.

58. Ribera A, Haurigot V, Garcia M, et al. Biochemical, histological and functional correction of mucopolysaccharidosis type IIIB by intra-cerebrospinal fluid gene therapy. *Hum. Mol. Genet.* Apr 01 2015;24(7):2078-2095.

59. Roca C, Motas S, Marco S, et al. Disease correction by AAV-mediated gene therapy in a new mouse model of mucopolysaccharidosis type IIID. *Hum. Mol. Genet.* Apr 15 2017;26(8):1535-1551.

60. Marco S, Haurigot V, Bosch F. In Vivo Gene Therapy for Mucopolysaccharidosis Type III (Sanfilippo Syndrome): A New Treatment Horizon. *Hum. Gene Ther.* Oct 2019;30(10):1211-1221.

- 61.** Agbandje-McKenna M, Kuhn R. Current opinion in virology: structural virology. *Curr. Opin. Virol.* Aug 2011;1(2):81-83.
- 62.** Mingozzi F, High KA. Immune responses to AAV in clinical trials. *Curr. Gene Ther.* Aug 2011;11(4):321-330.
- 63.** Naldini L. Gene therapy returns to centre stage. *Nature.* Oct 15 2015;526(7573):351-360.
- 64.** Buchlis G, Podsakoff GM, Radu A, et al. Factor IX expression in skeletal muscle of a severe hemophilia B patient 10 years after AAV-mediated gene transfer. *Blood.* Mar 29 2012;119(13):3038-3041.
- 65.** Hordeaux J, Hinderer C, Buza EL, et al. Safe and Sustained Expression of Human Iduronidase After Intrathecal Administration of Adeno-Associated Virus Serotype 9 in Infant Rhesus Monkeys. *Hum. Gene Ther.* Aug 2019;30(8):957-966.
- 66.** Jaen ML, Vila L, Elias I, et al. Long-Term Efficacy and Safety of Insulin and Glucokinase Gene Therapy for Diabetes: 8-Year Follow-Up in Dogs. *Molecular therapy. Methods & clinical development.* Sep 15 2017;6:1-7.
- 67.** Leone P, Shera D, McPhee SW, et al. Long-term follow-up after gene therapy for canavan disease. *Sci. Transl. Med.* Dec 19 2012;4(165):165ra163.
- 68.** Mittermeyer G, Christine CW, Rosenbluth KH, et al. Long-term evaluation of a phase 1 study of AADC gene therapy for Parkinson's disease. *Hum. Gene Ther.* Apr 2012;23(4):377-381.
- 69.** Sondhi D, Johnson L, Purpura K, et al. Long-term expression and safety of administration of AAVrh.10hCLN2 to the brain of rats and nonhuman primates for the treatment of late infantile neuronal ceroid lipofuscinosis. *Human gene therapy methods.* Oct 2012;23(5):324-335.
- 70.** Gilkes JA, Bloom MD, Heldermon CD. Preferred transduction with AAV8 and AAV9 via thalamic administration in the MPS IIIB model: A comparison of four rAAV serotypes. *Molecular genetics and metabolism reports.* Mar 2016;6:48-54.
- 71.** Fu H, Kang L, Jennings JS, et al. Significantly increased lifespan and improved behavioral performances by rAAV gene delivery in adult mucopolysaccharidosis IIIB mice. *Gene Ther.* Jul 2007;14(14):1065-1077.
- 72.** Murrey DA, Naughton BJ, Duncan FJ, et al. Feasibility and safety of systemic rAAV9-hNAGLU delivery for treating mucopolysaccharidosis IIIB: toxicology, biodistribution, and immunological assessments in primates. *Human gene therapy. Clinical development.* Jun 2014;25(2):72-84.
- 73.** Meadows AS, Duncan FJ, Camboni M, et al. A GLP-Compliant Toxicology and Biodistribution Study: Systemic Delivery of an rAAV9 Vector for the Treatment of Mucopolysaccharidosis IIIB. *Human gene therapy. Clinical development.* Dec 2015;26(4):228-242.

- 74.** Winner LK, Beard H, Hassiotis S, et al. A Preclinical Study Evaluating AAVrh10-Based Gene Therapy for Sanfilippo Syndrome. *Hum. Gene Ther.* May 2016;27(5):363-375.
- 75.** Falese L, Sandza K, Yates B, et al. Strategy to detect pre-existing immunity to AAV gene therapy. *Gene Ther.* Dec 2017;24(12):768-778.
- 76.** Boutin S, Monteilhet V, Veron P, et al. Prevalence of serum IgG and neutralizing factors against adeno-associated virus (AAV) types 1, 2, 5, 6, 8, and 9 in the healthy population: implications for gene therapy using AAV vectors. *Hum. Gene Ther.* Jun 2010;21(6):704-712.
- 77.** Calcedo R, Morizono H, Wang L, et al. Adeno-associated virus antibody profiles in newborns, children, and adolescents. *Clin. Vaccine Immunol.* Sep 2011;18(9):1586-1588.
- 78.** McIntyre C, Derrick-Roberts AL, Byers S, Anson DS. Correction of murine mucopolysaccharidosis type IIIA central nervous system pathology by intracerebroventricular lentiviral-mediated gene delivery. *J. Gene Med.* Nov-Dec 2014;16(11-12):374-387.
- 79.** Heldermon CD, Qin EY, Ohlemiller KK, et al. Disease correction by combined neonatal intracranial AAV and systemic lentiviral gene therapy in Sanfilippo Syndrome type B mice. *Gene Ther.* Sep 2013;20(9):913-921.
- 80.** Tardieu M, Zerah M, Husson B, et al. Intracerebral administration of adeno-associated viral vector serotype rh.10 carrying human SGSH and SUMF1 cDNAs in children with mucopolysaccharidosis type IIIA disease: results of a phase I/II trial. *Hum. Gene Ther.* Jun 2014;25(6):506-516.
- 81.** Tardieu M, Zerah M, Gougeon ML, et al. Intracerebral gene therapy in children with mucopolysaccharidosis type IIIB syndrome: an uncontrolled phase 1/2 clinical trial. *Lancet Neurol.* Sep 2017;16(9):712-720.
- 82.** Chen Y, Xu LP, Zhang XH, et al. Busulfan, Fludarabine, and Cyclophosphamide (BFC) conditioning allowed stable engraftment after haplo-identical allogeneic stem cell transplantation in children with adrenoleukodystrophy and mucopolysaccharidosis. *Bone Marrow Transplant.* Jun 2018;53(6):770-773.
- 83.** Taylor M, Khan S, Stapleton M, et al. Hematopoietic Stem Cell Transplantation for Mucopolysaccharidoses: Past, Present, and Future. *Biol. Blood Marrow Transplant.* Jul 2019;25(7):e226-e246.
- 84.** Tan EY, Boelens JJ, Jones SA, Wynn RF. Hematopoietic Stem Cell Transplantation in Inborn Errors of Metabolism. *Frontiers in pediatrics.* 2019;7:433.
- 85.** Sivakumur P, Wraith JE. Bone marrow transplantation in mucopolysaccharidosis type IIIA: a comparison of an early treated patient with his untreated sibling. *J. Inherit. Metab. Dis.* Oct 1999;22(7):849-850.
- 86.** Vellodi A, Young E, New M, Pot-Mees C, Hugh-Jones K. Bone marrow

transplantation for Sanfilippo disease type B. *J. Inherit. Metab. Dis.* 1992;15(6):911-918.

87. Aldenhoven M, Wynn RF, Orchard PJ, et al. Long-term outcome of Hurler syndrome patients after hematopoietic cell transplantation: an international multicenter study. *Blood.* Mar 26 2015;125(13):2164-2172.

88. Prasad VK, Mendizabal A, Parikh SH, et al. Unrelated donor umbilical cord blood transplantation for inherited metabolic disorders in 159 pediatric patients from a single center: influence of cellular composition of the graft on transplantation outcomes. *Blood.* Oct 1 2008;112(7):2979-2989.

89. Welling L, Marchal JP, van Hasselt P, van der Ploeg AT, Wijburg FA, Boelens JJ. Early Umbilical Cord Blood-Derived Stem Cell Transplantation Does Not Prevent Neurological Deterioration in Mucopolysaccharidosis Type III. *JIMD reports.* 2015;18:63-68.

90. Willing AE, Garbuzova-Davis SN, Zayko O, et al. Repeated administrations of human umbilical cord blood cells improve disease outcomes in a mouse model of Sanfilippo syndrome type III B. *Cell Transplant.* 2014;23(12):1613-1630.

91. Fraldi A, Serafini M, Sorrentino NC, Gentner B, Aiuti A, Bernardo ME. Gene therapy for mucopolysaccharidoses: in vivo and ex vivo approaches. *Ital. J. Pediatr.* Nov 16 2018;44(Suppl 2):130.

92. Ellison SM, Liao A, Wood S, et al. Pre-clinical Safety and Efficacy of Lentiviral Vector-Mediated Ex Vivo Stem Cell Gene Therapy for the Treatment of Mucopolysaccharidosis IIIA. *Molecular therapy. Methods & clinical development.* Jun 14 2019;13:399-413.

93. Holley RJ, Ellison SM, Fil D, et al. Macrophage enzyme and reduced inflammation drive brain correction of mucopolysaccharidosis IIIB by stem cell gene therapy. *Brain.* Jan 1 2018;141(1):99-116.

94. Langford-Smith A, Wilkinson FL, Langford-Smith KJ, et al. Hematopoietic stem cell and gene therapy corrects primary neuropathology and behavior in mucopolysaccharidosis IIIA mice. *Mol. Ther.* Aug 2012;20(8):1610-1621.

95. Sergijenko A, Langford-Smith A, Liao AY, et al. Myeloid/Microglial driven autologous hematopoietic stem cell gene therapy corrects a neuronopathic lysosomal disease. *Mol. Ther.* Oct 2013;21(10):1938-1949.

96. Clarke D, Pearse Y, Kan SH, et al. Genetically Corrected iPSC-Derived Neural Stem Cell Grafts Deliver Enzyme Replacement to Affect CNS Disease in Sanfilippo B Mice. *Molecular therapy. Methods & clinical development.* Sep 21 2018;10:113-127.

97. Espuny-Camacho I, Arranz AM, Fiers M, et al. Hallmarks of Alzheimer's Disease in Stem-Cell-Derived Human Neurons Transplanted into Mouse Brain. *Neuron.* Mar 8 2017;93(5):1066-1081 e1068.

98. Griffin TA, Anderson HC, Wolfe JH. Ex vivo gene therapy using patient iPSC-derived NSCs reverses pathology in the brain of a homologous mouse model. *Stem cell reports*. May 12 2015;4(5):835-846.

99. McGinley LM, Kashlan ON, Bruno ES, et al. Human neural stem cell transplantation improves cognition in a murine model of Alzheimer's disease. *Sci. Rep.* Oct 3 2018;8(1):14776.

100. Snyder EY, Taylor RM, Wolfe JH. Neural progenitor cell engraftment corrects lysosomal storage throughout the MPS VII mouse brain. *Nature*. Mar 23 1995;374(6520):367-370.

101. Zhang T, Ke W, Zhou X, et al. Human Neural Stem Cells Reinforce Hippocampal Synaptic Network and Rescue Cognitive Deficits in a Mouse Model of Alzheimer's Disease. *Stem cell reports*.