

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,000

Open access books available

116,000

International authors and editors

120M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Next-Generation Therapeutics: mRNA as a Novel Therapeutic Option for Single-Gene Disorders

Tatjana Michel, Hans-Peter Wendel and
Stefanie Krajewski

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/62243>

Abstract

In single-gene disorders, such as α 1-antitrypsin deficiency (AATD), hemophilia B (clotting factor IX deficiency), and lecithin-cholesterol acyltransferase deficiency (LCATD), a gene mutation causes missing or dysfunctional protein synthesis, which in turn can lead to serious complications for the patient affected. Furthermore, single-gene disorders are associated with severe, early-onset conditions and necessitate expensive lifelong care. Today, therapeutic treatment options remain limited, cost-intensive, or ineffective. Therefore, the novel mRNA-based therapeutic strategy for the treatment of single-gene disorders, which is based on the induction of *de novo* synthesis of the functional proteins, has extraordinary potential. After the delivery of the specific mRNA to the target cells, the desired protein is expressed by the cells' own translational machinery, and hence, a fully functional protein replaces the defective or missing protein. mRNA therapy provides an innovative, highly promising, and inexpensive therapeutic approach and will thus lead to new advances in the treatment of single-gene disorders.

Keywords: α 1-antitrypsin deficiency, hemophilia B, lecithin-cholesterol acyltransferase deficiency, mRNA therapy, single-gene disorders, messenger RNA, next-generation therapeutics, gene mutation

1. Introduction

The human body is made up of millions of cells, which have special, well-defined functions, such as the transportation of oxygen in the blood. Proteins carry out all these functions that are necessary for life. In most of the cells, the genetic information is encoded on the 23 pairs of

chromosomes found in the nucleus. On each chromosome, the information for the production of a great range of different proteins is contained in genes made up of DNA. There are approximately 25,000 protein-encoding genes in the human genome [1].

Protein synthesis takes place in two major processes. First, the DNA is transcribed into the mRNA in the nucleus followed by cytoplasmic mRNA translation into the protein.

Alteration in a single gene (i.e., a single-gene disorder) caused by a mutation in the gene's DNA sequences leads to dysfunction of the gene. As a result, the protein the gene codes for is either altered or missing, which can result in serious complications in the human body.

Over the last few years, it has become apparent that single-gene disorders are far more numerous than previously assumed, and more than 1800 single-gene disorders have been identified [2]. Moreover, single-gene disorders are associated with severe, early-onset conditions necessitating lifelong care [3].

2. Examples of single-gene disorders

2.1. α 1-Antitrypsin (AAT) deficiency (AATD)

AAT is a serine protease inhibitor belonging to the serpin superfamily, and it is predominantly synthesized in the liver and released into the bloodstream. By inhibiting neutrophil elastase, which is released from activated neutrophil granulocytes and macrophages, AAT plays a pivotal role in the prevention of proteolytic damage to the host tissue in the presence of inflammatory processes. Physiological AAT plasma concentrations are within the range of 20 to 53 $\mu\text{mol/L}$ (150–300 mg/dL).

Mutations in the SERPIN1A gene are associated with AATD, one of the most common metabolic diseases in Europe. AATD, first described in 1963 by Laurell and Erikson, affects approximately 1 in 2000 to 1 in 5000 individuals [4, 5]. People suffering from AATD have AAT levels below 11 $\mu\text{mol/L}$ (80 mg/dL) and are predisposed to severe lung diseases, such as chronic obstructive pulmonary disease (COPD), or liver diseases.

2.2. Hemophilia B

Hemophilia B, also called Christmas disease, is the second most common form of hemophilia, affecting approximately 20% of those diagnosed with hemophilia. It is a bleeding disorder that is typically inherited and characterized by a lack of clotting activity of factor IX (FIX; Christmas factor), a blood clotting factor that is synthesized in hepatocytes and plays a crucial role in blood coagulation. Depending on the bleeding phenotype of hemophilia B, which is classified as mild, moderate, or severe, and the sufferer's overall health, symptoms vary from prolonged, partially excessive bleeding to serious bruising and joint pain [6].

2.3. Lecithin-cholesterol acyltransferase (LCAT) deficiency (LCATD)

The gene for LCAT is localized on chromosome 16 and primarily expressed in the liver. After secretion to the plasma, it is primarily found attached to circulating high-density lipoprotein (HDL) particles [7]. LCAT converts free cholesterol into cholesterol esters on the surface of HDL, thus removing cholesterol from the blood and tissues [8].

Mutations in the LCAT gene result in deficient or absent catalytic LCAT activity, leading to a reduction in the enzyme's ability to attach cholesterol to lipoproteins. Hence, deficiency leads to the accumulation of unesterified cholesterol in different tissues (e.g., in the cornea, erythrocytes, or kidneys) and may lead to corneal opacities, renal failure, or hemolytic anemia [9]. Due to the accumulation of cholesterol in the lining of the arteries, LCATD sufferers have an increased risk for premature atherosclerosis.

2.4. Familial hypercholesterolemia (FH)

FH is a genetic disease characterized by high low-density lipoprotein (LDL) cholesterol levels in the blood. This is caused by a defect in the gene for the LDL receptor (LDLR) that prevents it from absorbing the LDL from the blood into the cell for metabolism. The symptoms in patients with FH range from harmless fatty skin deposits called xanthoma to life-threatening atherosclerotic vascular disease, which can culminate in myocardial infarction or stroke [10]. In patients suffering from the homozygous form of FH, such life-threatening complications occur in infancy. Statins are currently used for standard therapy, but their efficacy is controversial [11].

2.5. Available treatment options

Different treatment options exist depending on the genetic disorder. Treatment of AATD is currently achieved by aerosol or intravenous augmentation therapy of purified and pooled human plasma AAT protein [12]. Furthermore, the risks of proinflammatory stimuli to the lung need to be minimized by ensuring that the patient abstains from smoking, using bronchodilators, etc. Augmentation therapy is, however, very expensive, because the AAT protein derived from a healthy donor needs to undergo a rigorous screening process before it is ready to be infused [13].

Patients suffering from hemophilia B are routinely treated with recombinant FIX concentrates, which have greatly reduced the mortality associated with hemophilia B. However, there are still significant drawbacks of this existing therapy, including the necessity of multiple weekly infusions for patients with severe hemophilia B as well as repeated bleeding despite prophylactic therapy, which can cause long-term damage in joints and other tissues.

The therapeutic up-regulation of LCAT function has gained interest in recent years, not only as an enzyme replacement therapy for LCATD syndromes but also as a potential therapeutic strategy for reducing atherosclerosis [14]. In 2013, the first case report was published highlighting the success of LCAT replacement therapy using a recombinant enzyme form in a 53-year-old patient.

On the contrary, many gene therapy-based systems for the treatment of various genetic disorders have been developed and investigated during the last few years. Gene therapy promises the permanent expression of the functional protein after the incorporation of the corresponding gene into the host genome. However, as yet, gene therapy has not found wide clinical application, because, depending on the vectors used, it can be associated with risks for the patient, such as insertional mutagenesis, carcinogenic effects, immune responses, low gene-transfer efficiency, or protein misfolding [15].

3. mRNA as a novel therapeutic option

3.1. mRNA as a therapeutic agent

Next to gene therapy, wherein genetic defects are corrected by the introduction of specific DNA sequences into the genome, mRNA-based therapy promises new advances in the treatment of single-gene disorders. Via the delivery of a specific *in vitro*-generated mRNA to the target cells, the expression of a desired protein can be induced. The idea of using specific mRNAs to produce a protein of interest instead of protein replacement via DNA gene therapy was described 25 years ago by Wolff and colleagues [16]. At that time, however, the stability of the mRNA was poor and the immunogenicity was too high for the therapy to be practical. In recent years, the administration of the mRNA as a therapeutic agent has gained enormous potential in the fields of disease treatment, regenerative medicine, and vaccination [17–21].

Especially for monogenic diseases, mRNA therapy can be a highly beneficial alternative to classical gene therapy. Because monogenic diseases result in defective or missing protein synthesis, protein replacement therapy is primarily used for these kinds of diseases. The mRNA can be easily produced in large amounts for the protein of interest in comparison to pDNA, and if the mRNA is used, there is no need to integrate promoter and terminator regions in the sequence [22].

Moreover, the mRNA-based therapeutic strategy has significant advantages:

1. The introduced mRNA is translated by the cell's own translational machinery under physiological conditions.
2. By introducing a specific mRNA, the physiological state of the cell is not altered, because the effect is transient and not mutagenic.
3. The protein synthesis can be controlled directly without intervening in the human genome.
4. The mRNA does not need to enter the nucleus for translation [15, 20, 22].

Overall, this therapeutic strategy could be safely used in patients, and it is more cost-effective and easier to manipulate than gene therapy.

The standard *in vitro* procedure for mRNA generation begins with the plasmid, which contains the coding sequence of the protein. First, this sequence is amplified using polymerase chain

reaction (PCR). The generated DNA template contains all the important elements of the mRNA. Second, the amplified PCR product is used to generate the mRNA. Therefore, *in vitro* transcription (IVT) is performed using the T7 or SP6 RNA polymerase, which synthesizes the mRNA. After the purification and quality control steps, the mRNA is ready to use (Figure 1).




Figure 1. Generation process of the mRNA.

The mRNA is a single-stranded molecule containing a poly(A)-tail at the end and a 5'-cap at the beginning. The coding sequence for the protein is marked by a start codon and a stop codon. The untranslated regions (UTRs) are in between the cap/tail and the coding sequence (Figure 2) [16].



Figure 2. Schematic overview of the general structure of an IVT mRNA.

However, IVT mRNA is very sensitive to degradation by nucleases, which limits its suitability for transfections and therapeutic applications [21]. If the mRNA will be applied as a therapeutic agent, there is a need for special modifications. The modifications should be nonmutagenic and should not interfere with the translation machinery of the cell [23]. To improve the translation of the mRNA, the cap and the poly(A)-tail are important. The mRNA cap is responsible for the recognition of the ribosomes and binding to the ribosomes and for the initiation of the translation machinery [24]. It was reported that part of the unmodified cap structure could be bound in the wrong direction to the mRNA during IVT. Thus, the mRNA cannot be located by the cap-binding complex (CBC) of the ribosomes and the translation cannot follow [25]. By using an anti-reverse cap analog (ARCA) during IVT, this can be prevented. An ARCA contains a 5'-5' triphosphate bridge, and the 3'OH is replaced by OCH₃, making it impossible for the cap to bind in the wrong direction [26]. The 5'-5' bridge and the chemical modifications at the 3'-position lead to translatable mRNA of high quality and great translation efficiency [27, 28]. Furthermore, ARCA-capped mRNA is more resistant

to hydrolases [29]. The poly(A)-tail, which binds to the polyadenosyl-binding-protein (PABP), is also very important for the stability and translation of the mRNA [30, 31]. The PABP interacts with the CBC and makes a circular structure of the mRNA molecule [15, 32]. This circular structure minimizes the contact surface for nucleases [33]. If the poly(A)-tail is shorter than 12 adenine nucleosides or removed completely, the cap structure is cleaved and the mRNA is degraded. Thus, the poly(A)-tail is very important to obtaining the cap structure and delaying the degradation of the mRNA [15, 34]. A poly(A)-tail of more than 60 adenine monophosphates increases the translation efficiency of the mRNA [35]. Therefore, the poly(A)-tail and the cap structure contribute to the stability of the mRNA [27, 36]. The 5'- and 3'-UTRs include specific regulatory sequence regions that are necessary to modulate the stability and translation of the mRNA [16]. The 3'-UTR region contains α - and β -globin sequences that enhance the stability and translation of the mRNA [16, 37, 38]. Furthermore, the 3'- and 5'-UTR regions inhibit the decapping and degradation of the mRNA [16, 39, 40]. On the contrary, if limited protein production is desired, faster mRNA degradation is possible by integrating AU-rich areas in the 3'-UTR region [41]. The protein-coding region of the mRNA can be optimized in two ways. First, the use of optimized codons leads to improved translation of the sequence to the desired protein. Second, via the optimization of the bases, the endonucleolytic degradation can be reduced [16].

3.2. Potential immune response against IVT mRNA

In comparison to recombinant protein in the substitution therapy approach, mRNA-translated proteins are autologously produced by the cell's own machinery. Usually, they undergo the correct modification and folding [16], but protein aggregation and immune reactions to the translated protein cannot be excluded. Therefore, it is necessary to investigate the activation of the immune system and distribution of the antibodies against IVT mRNA-translated proteins in clinical studies. Furthermore, the combination of the immune reaction to the translated mRNA protein and the foreign mRNA within the cells can lead to immunopathology in the cells or organs [16, 42].

Regarding protein replacement therapies, the activation of the immune system caused by mRNA administration might be disadvantageous. It is well known that foreign mRNAs as well as pDNAs activate the immune system via recognition through toll-like receptors (TLRs). TLRs recognize different pathogen patterns resulting in the expression of different cytokines. TLR3 and TLR7/8 are responsible for the recognition of different RNA types [43]. The mRNA and other RNA types, such as small interfering RNA (siRNA) or double-stranded RNA (dsRNA), are recognized by TLR3 [44–46], whereas TLR7/8 is activated by single-stranded RNA (ssRNA) [47]. In nonimmune cells, the recognition of the mRNA occurs through the retinoic acid-inducible gene I (RIG-I), which is activated by short RNA and dsRNA [48] and leads to interleukin-1 β (IL-1 β) production [49]. It is known that the IVT mRNA leads to a strong distribution of tumor necrosis factor- α (TNF- α) if the mRNA has no modifications [50]. Additionally, a strong type I interferon (type I IFN) response of the cells is induced upon contact with exogenous mRNAs. This is also induced by mRNAs that form secondary structures, such as loops or hairpins, or by mRNAs that bind to incompletely synthesized

mRNA fragments or incompletely degraded DNA fragments [43, 51]. IFNs then activate the antiviral genes in the genome and lead to a translation stop and the degradation of the mRNA [43]. However, the immune reaction can be avoided if the mRNA is purified and if modified nucleotides are inserted during IVT [50, 52–54].

The incorporation of different modified nucleotides, such as pseudouridine (pseudo-U), 2'-thiouridine (2'-tU), 5'-methyluridine (5'-mU), 5-methylcytosine (5'-mC), or N⁶-methyladenosine (N⁶-mA), can prevent the cellular immune response [50, 53]. These modified nucleotides incorporated in the mRNA help to avoid the activation of TLRs [50]. In particular, pseudo-U and 2'-tU make the recognition of IVT mRNA by RIG-I impossible [16, 54, 55]. To minimize the immune activation and optimize the translation efficiency of the mRNA to the protein, high-performance liquid chromatography (HPLC) purification should be applied to eliminate the dsRNA contamination that can still be present after the IVT process [56].

However, mRNA modification may also represent another risk. The naturally existing mRNA is degradable by the RNases in the extracellular space, but the modification of the IVT mRNA makes degradation more difficult [57]. Some of the modified nucleotides are associated with mitochondrial toxicities and hepatic failure and play a role in viral and tumor cell replication [58, 59]. Here too, further investigations and clinical trials are necessary to prove the risks and benefits of IVT mRNA modifications.

3.3. mRNA delivery systems and specific targeting

For clinical mRNA applications, a specific delivery system is needed, because otherwise the delivery of the mRNA to the target cells is unguaranteed and inefficient [27]. Therefore, the development and engineering of safe and effective delivery vectors for mRNA therapy is inevitable [60]. Viral and nonviral vectors can be used to bring the mRNA into the cells. For the direct translation of the mRNA to the protein, only positive-stranded viruses can be applied [61]. Negative-stranded viruses are not infectious and need an RNA-dependent RNA polymerase for mRNA translation [62]. However, viral vectors have some limitations; for example, they can be carcinogenic [63], they can activate immune responses [64], they can be difficult to produce [65], and they have a limited packaging capacity [66].

Nonviral vectors have lower immunogenicity, they can deliver large genetic molecules, and they are easier to produce [60, 67, 68]. Nonviral delivery systems can be subdivided into direct and indirect delivery systems. Direct delivery is possible via electroporation or gene guns. Electroporation is an early and efficient method of transporting mRNA into cells [69], whereby electrical pulses make the cell membrane permeable for the entry of the mRNA into the cytosol. This method does not induce immune activation, which may occur when mRNA carriers are used [27, 69]. Transfection using a gene gun requires heavy metal particles to get the nucleic acid into the cell [70]. This method allows the delivery of the mRNA to mammalian organs with minimal damage and leads to transient protein expression in the target tissues [70]. Self-assembled complexation of negatively charged mRNAs and positively charged liposomes and polysomes to lipoplexes or polyplexes is the most widely used method of bringing mRNA into cells [60]. The lipoplexes can be taken up by the cells in two different ways. The first way is by the endocytosis of the lipoplexes, whereby 98% of the lipoplexes enter the cell. The second way

is through the fusion of the cell membrane and the lipoplexes, which results in the uptake of the remaining mRNA [71]. After the release of the mRNA into the cytosol, the protein translation can begin. The encapsulation of the mRNA into liposomes is a rapid, transient, and cell cycle-independent delivery method [27]. The translation of the mRNA to the protein can be measured 1 h after the transfection in nondividing cells [72].

For *in vivo* applications, the perfect mRNA delivery vector has to overcome various barriers:

1. Protection against nucleases,
2. Avoidance of nonspecific interactions with proteins or cells,
3. Prevention of renal clearance,
4. Permission of extravasation to target the tissue of interest, and
5. Increased entry into the cell [60].

For systemic delivery, the lipid and polymer complexes show protective properties against nucleases, whereby the mRNA is protected and the stability is increased [73]. However, liposome complexes sometimes interact with serum proteins. Together, they form aggregates or clots and are cleared rapidly [74]. The conjugation of the complexes with polyethylene glycol (PEG) helps to inhibit the nonspecific uptake and attachment to serum proteins [75].

Nanoscale delivery systems (10–200 nm) enhance the uptake efficiency and reduce the systemic toxicity [75]. The use of PEG coating of the liposomes increases the blood circulation time and avoids the detection of liposomes by immune cells [75, 76]. The liposomes have many advantages, such as low batch-to-batch variability, easy synthesis, biocompatibility, and scalability, over many other delivery systems [75]. The liposome surface can be functionalized by conjugation to chemically reactive lipid head groups [77]. This property makes it possible to functionalize the surface with ligands and thus enhance the target delivery [75, 78].

Different methods of application are tested for *in vivo* delivery. Polyplex nanomicelles applied with hydrodynamic intravenous injection have been shown to effectively deliver the mRNA to the liver in mouse models. This method shows a strong protein expression in nearly 100% of liver cells [79]. Intramuscular or intraperitoneal injection of erythropoietin (EPO)-coding mRNA complexed with cationic lipids leads to significantly high levels of EPO *in vivo* [20, 80]. Intratracheal and intranasal applications of Foxp3 mRNA show protective properties against asthma in mice [81]. Furthermore, it has been shown that intradermal [82] and intranodal [83] applications of the mRNA in animal models resulted in immunization against tumors. Many other methods of applying the mRNA have been described in publications in recent years, and this field is developing rapidly [19, 84]. Furthermore, patient-centered applications have improved, especially for mRNA therapy. Nebulization with the Pari-Boy® is the standard method used to apply the drugs to the lung. A study on the influence on mRNA transfection efficiency shows that the nebulization of complexed mRNA has no effects *in vitro* [85].

3.4. mRNA applications

Gene therapy allows the replacement of a defective gene through substitution and integration of the correct genetic code in the genome. The genome integration of this genetic code via viruses guarantees highly efficient gene replacement methods. However, undesirable effects, such as mutagenesis and innate immune response, may jeopardize the life or safety of the patient [27, 86]. Nonviral gene delivery is safer, but it is also associated with lower transfection efficiency due to insufficient nuclear transport. Furthermore, modifications of the pDNA, including adding a strong constructive promoter to improve transcription efficiency, may lead to unexpected alterations in the genome [27, 86, 87].

Protein-substitution therapy is associated with adverse reactions, such as headache, dizziness, and nausea [88], and high costs [89]. Recombinant proteins have been expressed in different microorganisms [90–92], plant cells [93], and human cells [94–96], but they are linked to disadvantages such as nonglycosylation or incorrect glycosylation of the product as well as product contaminations with endotoxins. Plasma-derived proteins, which are purified from human donor blood, are limited and prone to contamination.

The mRNA is an alternative to overcome the disadvantages of pDNA and direct protein substitution [15]. The mRNA does not need to enter the nucleus for transcription, because the mRNA is directly translated in the cytosol of the cells; thus, the insertion of exogenous DNA into the genome poses no risk. Furthermore, the mRNA uses the cell's own translation machinery and requires no strong promoter. Effective mRNA transfer can also take place in nondividing cells, and immunogenicity can be overcome by different modifications of the mRNA molecule [27]. Moreover, compared to protein-substitution therapy, the expression of receptors and intracellular molecules can be induced with specific mRNAs (**Figure 3**).




Figure 3. The way from encapsulated mRNA to protein expression in the target cell.

Several studies describe that the use of different mRNAs leads to an increase in the respective protein *in vitro* and *in vivo*. The first application of the mRNA was performed in 1992 [97]. However, the broad application of IVT mRNA only became possible as modified nucleotides were used; thus, a reduction of immune reactions and an increase of mRNA translation efficiency were achieved [50, 56]. For example, the successful expression of surfactant protein B (SP-B) of therapeutically relevant levels was shown in a mouse model after the application of SP-B-encoding mRNA via direct administration to the lung. The results showed that the inflammation of the lung (and the inevitable respiratory failure and death) was prevented [20, 98]. Likewise, the expression of the regulatory T-cell transcription factor (FOXP3) led to the prevention of asthma in a mouse model. After the administration of FOXP3-coding mRNA to the lung, the expressed protein protected the lung from allergen-induced inflammation. This mRNA approach can be used as a preventive and therapeutic drug [81]. In a different study, the expression of the AAT protein was shown after the transfection of cells with AAT-encoding mRNA. The level of AAT was measurable in the cellular supernatant 48 h after transfection, and the functionality of the protein was proven. All these studies show that mRNA therapy also promises a novel therapeutic strategy in the treatment of single-gene diseases such as AATD [99].

In the case of FH, it is also possible to induce the expression of functional LDLR after the transfection and thus regulate LDL metabolism. In this case, functional LDLR expression can help to prevent secondary diseases, such as stroke and atherosclerotic plaques. Although the induction of the expression of proteins, such as LCAT and FIX, can be performed similarly to the approach of AATD-mRNA therapy, different organs are targeted. Regarding hemophilia B and LCATD, hepatocytes should be transfected, which can be achieved by intravenous injections of the complexed protein-encoding mRNA.

Depending on the application method and the genetic defect, it is important that the cell type of interest is targeted and transfected with IVT mRNA. Additionally, the dose-effect relationship could be a challenge *in vivo*, whereby the bioavailability and individual variations also play a pivotal role and potentially make individual dose adaptation necessary [16]. Overall, mRNA application promises an effective and low-cost therapeutic strategy with the potential to efficiently correct serious monogenic diseases.

3.5. Summary

Overall, the mRNA as a novel therapy for monogenic diseases has many advantages over the currently existing treatment options such as substitution therapy or gene therapy. Using the mRNA, the expression of nearly all proteins can be induced. In this way, not only extracellular proteins, but also proteins, which are important for the function of the cells, such as growth factors or receptors can be generated. Furthermore, the mRNA does not need to penetrate into the nucleus. The effect of the mRNA is transient, thereby enabling the precise control of protein expression. Depending on the introduced modifications, the half-life of the respective mRNA and thus a reduction or increase in mRNA stability can be determined. Developments in recent years, such as modification and purification methods, have made it possible to use the mRNA as a therapeutic agent, because it became possible to control the immune reaction, which is

typically triggered by exogenous mRNA. Further developments increased the extracellular stability and made mRNA transfection of the cells with nonviral vectors possible and efficient. The mRNA can be produced in large batches and under good manufacturing practice (GMP) conditions without batch-to-batch variations. Thereby, the production of specific mRNA is significantly cheaper compared to the production of the corresponding protein for substitution therapy.

The mRNA as a therapeutic agent could be a great help for patients suffering from monogenic diseases. The flexibility and variability of proteins that can be replaced by the cell's own translational machinery through the use of the mRNA is nearly unlimited. This makes mRNA to a unique therapeutic molecule, which will revolutionize therapeutic options for affected patients in the coming years.

Author details

Tatjana Michel, Hans-Peter Wendel and Stefanie Krajewski*

*Address all correspondence to: stefanie.krajewski@uni-tuebingen.de

Department of Thoracic, Cardiac and Vascular Surgery, University of Tuebingen, Tuebingen, Germany

References

- [1] Collins FS, Lander ES, Rogers J, Waterston RH, Conso IHGS: Finishing the euchromatic sequence of the human genome. *Nature* 2004, 431(7011):931–945.
- [2] Antonarakis SE, Beckmann JS: Mendelian disorders deserve more attention. *Nat Rev Genet* 2006, 7(4):277–282.
- [3] Ropers HH: Single gene disorders come into focus—again. *Dialogues Clin Neurosci* 2010, 12(1):95–102.
- [4] Laurell CB, Eriksson S: The electrophoretic α 1-globulin pattern of serum in α 1-antitrypsin deficiency. 1963. *COPD* 2013, 10 Suppl 1:3–8.
- [5] Hurley K, Lacey N, O'Dwyer CA, Bergin DA, McElvaney OJ, O'Brien ME, McElvaney OF, Reeves EP, McElvaney NG: α -1 antitrypsin augmentation therapy corrects accelerated neutrophil apoptosis in deficient individuals. *J Immunol* 2014, 193(8):3978–3991.
- [6] Cancio MI, Reiss UM, Nathwani AC, Davidoff AM, Gray JT: Developments in the treatment of hemophilia B: focus on emerging gene therapy. *Appl Clin Genet* 2013, 6:91–101.

- [7] Francone OL, Gurakar A, Fielding C: Distribution and functions of lecithin: cholesterol acyltransferase and cholesteryl ester transfer protein in plasma lipoproteins. Evidence for a functional unit containing these activities together with apolipoproteins A-I and D that catalyzes the esterification and transfer of cell-derived cholesterol. *J Biol Chem* 1989, 264(12):7066–7072.
- [8] Kuivenhoven JA, Pritchard H, Hill J, Frohlich J, Assmann G, Kastelein J: The molecular pathology of lecithin:cholesterol acyltransferase (LCAT) deficiency syndromes. *J Lipid Res* 1997, 38(2):191–205.
- [9] McIntyre N: Familial LCAT deficiency and fish-eye disease. *J Inherited Metab Dis* 1988, 11(Suppl 1):45–56.
- [10] Raal FJ, Santos RD: Homozygous familial hypercholesterolemia: current perspectives on diagnosis and treatment. *Atherosclerosis* 2012, 223(2):262–268.
- [11] Pijlman AH, Huijgen R, Verhagen SN, Imholz BP, Liem AH, Kastelein JJ, Abbink EJ, Stalenhoef AF, Visseren FL: Evaluation of cholesterol lowering treatment of patients with familial hypercholesterolemia: a large cross-sectional study in The Netherlands. *Atherosclerosis* 2010, 209(1):189–194.
- [12] American Thoracic Society, European Respiratory Society: American Thoracic Society/ European Respiratory Society statement: standards for the diagnosis and management of individuals with α -1 antitrypsin deficiency. *Am J Respir Crit Care Med* 2003, 168(7): 818–900.
- [13] Petrache I, Hajjar J, Campos M: Safety and efficacy of α -1-antitrypsin augmentation therapy in the treatment of patients with α -1-antitrypsin deficiency. *Biol Targets Ther* 2009, 3:193–204.
- [14] Kunnen S, Van Eck M: Lecithin:cholesterol acyltransferase: old friend or foe in atherosclerosis? *J Lipid Res* 2012, 53(9):1783–1799.
- [15] Yamamoto A, Kormann M, Rosenecker J, Rudolph C: Current prospects for mRNA gene delivery. *Eur J Pharm Biopharm* 2009, 71(3):484–489.
- [16] Sahin U, Kariko K, Tureci O: mRNA-based therapeutics—developing a new class of drugs. *Nat Rev Drug Discov* 2014, 13(10):759–780.
- [17] Kübler H, Maurer T, Stenzl A, Feyerabend S, Steiner U, Schostak M, Schultze-Seemann W, vom Dorp F, Pilla L, Viatali G, et al.: Final analysis of a phase I/IIa study with CV9103, an intradermally administered prostate cancer immunotherapy based on self-adjuvanted mRNA. *J Clin Oncol* 2011, 29:498–499.
- [18] Fotin-Mleczek M, Duchardt KM, Lorenz C, Pfeiffer R, Ojkic-Zrna S, Probst J, Kallen KJ: Messenger RNA-based vaccines with dual activity induce balanced TLR-7 dependent adaptive immune responses and provide antitumor activity. *J Immunother* 2011, 34(1): 1–15.

- [19] Petsch B, Schnee M, Vogel AB, Lange E, Hoffmann B, Voss D, Schlake T, Thess A, Kallen KJ, Stitz L, et al.: Protective efficacy of *in vitro* synthesized, specific mRNA vaccines against influenza A virus infection. *Nat Biotechnol* 2012, 30(12):1210–1216.
- [20] Kormann MS, Hasenpusch G, Aneja MK, Nica G, Flemmer AW, Herber-Jonat S, Huppmann M, Mays LE, Illenyi M, Schams A, et al.: Expression of therapeutic proteins after delivery of chemically modified mRNA in mice. *Nat Biotechnol* 2011, 29(2):154–157.
- [21] Avci-Adali M, Behring A, Keller T, Krajewski S, Schlensak C, Wendel HP: Optimized conditions for successful transfection of human endothelial cells with *in vitro* synthesized and modified mRNA for induction of protein expression. *J Biol Eng* 2014, 8(1):8.
- [22] Tavernier G, Andries O, Demeester J, Sanders NN, De Smedt SC, Rejman J: mRNA as gene therapeutic: how to control protein expression. *J Control Release* 2011, 150(3):238–247.
- [23] Wang X, He C: Dynamic RNA modifications in posttranscriptional regulation. *Mol Cell* 2014, 56(1):5–12.
- [24] Gallie DR: The cap and poly(A) tail function synergistically to regulate mRNA translational efficiency. *Genes Dev* 1991, 5(11):2108–2116.
- [25] Matsuo H, Li H, McGuire AM, Fletcher CM, Gingras AC, Sonenberg N, Wagner G: Structure of translation factor eIF4E bound to m7GDP and interaction with 4E-binding protein. *Nat Struct Biol* 1997, 4(9):717–724.
- [26] Stepinski J, Waddell C, Stolarski R, Darzynkiewicz E, Rhoads RE: Synthesis and properties of mRNAs containing the novel “anti-reverse” cap analogs 7-methyl(3'-O-methyl)GpppG and 7-methyl (3'-deoxy)GpppG. *RNA* 2001, 7(10):1486–1495.
- [27] Youn H, Chung JK: Modified mRNA as an alternative to plasmid DNA (pDNA) for transcript replacement and vaccination therapy. *Expert Opin Biol Ther* 2015, 15(9):1337–1348.
- [28] Mockey M, Goncalves C, Dupuy FP, Lemoine FM, Pichon C, Midoux P: mRNA transfection of dendritic cells: synergistic effect of ARCA mRNA capping with poly(A) chains in cis and in trans for a high protein expression level. *Biochem Biophys Res Commun* 2006, 340(4):1062–1068.
- [29] Grudzien E, Kalek M, Jemielity J, Darzynkiewicz E, Rhoads RE: Differential inhibition of mRNA degradation pathways by novel cap analogs. *J Biol Chem* 2006, 281(4):1857–1867.
- [30] Chang H, Lim J, Ha M, Kim VN: TAIL-seq: genome-wide determination of poly(A) tail length and 3' end modifications. *Mol Cell* 2014, 53(6):1044–1052.
- [31] Peng J, Murray EL, Schoenberg DR: *In vivo* and *in vitro* analysis of poly(A) length effects on mRNA translation. *Methods Mol Biol* 2008, 419:215–230.

- [32] Gallie DR: A tale of two termini: a functional interaction between the termini of an mRNA is a prerequisite for efficient translation initiation. *Gene* 1998, 216(1):1–11.
- [33] Newbury SF: Control of mRNA stability in eukaryotes. *Biochem Soc Trans* 2006, 34(Pt 1):30–34.
- [34] Schwartz D, Decker CJ, Parker R: The enhancer of decapping proteins, Edc1p and Edc2p, bind RNA and stimulate the activity of the decapping enzyme. *RNA* 2003, 9(2): 239–251.
- [35] Elango N, Elango S, Shivshankar P, Katz MS: Optimized transfection of mRNA transcribed from a d(A/T)₁₀₀ tail-containing vector. *Biochem Biophys Res Commun* 2005, 330(3):958–966.
- [36] Parker R, Sheth U: P bodies and the control of mRNA translation and degradation. *Mol Cell* 2007, 25(5):635–646.
- [37] Kariko K, Kuo A, Barnathan E: Overexpression of urokinase receptor in mammalian cells following administration of the *in vitro* transcribed encoding mRNA. *Gene Ther* 1999, 6(6):1092–1100.
- [38] Ross J, Sullivan TD: Half-lives of β and γ globin messenger RNAs and of protein synthetic capacity in cultured human reticulocytes. *Blood* 1985, 66(5):1149–1154.
- [39] Kuhn AN, Beibetaert T, Simon P, Vallazza B, Buck J, Davies BP, Tureci O, Sahin U: mRNA as a versatile tool for exogenous protein expression. *Curr Gene Ther* 2012, 12(5): 347–361.
- [40] Bergman N, Moraes KC, Anderson JR, Zaric B, Kambach C, Schneider RJ, Wilusz CJ, Wilusz J: Lsm proteins bind and stabilize RNAs containing 5' poly(A) tracts. *Nat Struct Mol Biol* 2007, 14(9):824–831.
- [41] Chen CY, Shyu AB: AU-rich elements: characterization and importance in mRNA degradation. *Trends Biochem Sci* 1995, 20(11):465–470.
- [42] Racanelli V, Rehmann B: The liver as an immunological organ. *Hepatology* 2006, 43(2 Suppl 1):S54–S62.
- [43] Janeway C: Immunologie, 7. Aufl. edn. Heidelberg/Berlin: Spektrum, Akadem. Verl.; 2009.
- [44] Kariko K, Ni H, Capodici J, Lamphier M, Weissman D: mRNA is an endogenous ligand for Toll-like receptor 3. *J Biol Chem* 2004, 279(13):12542–12550.
- [45] Alexopoulou L, Holt AC, Medzhitov R, Flavell RA: Recognition of double-stranded RNA and activation of NF- κ B by Toll-like receptor 3. *Nature* 2001, 413(6857):732–738.
- [46] Kariko K, Bhuyan P, Capodici J, Weissman D: Small interfering RNAs mediate sequence-independent gene suppression and induce immune activation by signaling through toll-like receptor 3. *J Immunol* 2004, 172(11):6545–6549.

- [47] Heil F, Hemmi H, Hochrein H, Ampenberger F, Kirschning C, Akira S, Lipford G, Wagner H, Bauer S: Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8. *Science* 2004, 303(5663):1526–1529.
- [48] Schlee M, Roth A, Hornung V, Hagmann CA, Wimmenauer V, Barchet W, Coch C, Janke M, Mihailovic A, Wardle G, et al.: Recognition of 5' triphosphate by RIG-I helicase requires short blunt double-stranded RNA as contained in panhandle of negative-strand virus. *Immunity* 2009, 31(1):25–34.
- [49] Poeck H, Bscheider M, Gross O, Finger K, Roth S, Rebsamen M, Hanneschlagler N, Schlee M, Rothenfusser S, Barchet W, et al.: Recognition of RNA virus by RIG-I results in activation of CARD9 and inflammasome signaling for interleukin 1 β production. *Nat Immunol* 2010, 11(1):63–69.
- [50] Kariko K, Buckstein M, Ni H, Weissman D: Suppression of RNA recognition by Toll-like receptors: the impact of nucleoside modification and the evolutionary origin of RNA. *Immunity* 2005, 23(2):165–175.
- [51] Rautsi O, Lehmusvaara S, Salonen T, Hakkinen K, Sillanpaa M, Hakkarainen T, Heikkinen S, Vahakangas E, Yla-Herttuala S, Hinkkanen A, et al.: Type I interferon response against viral and non-viral gene transfer in human tumor and primary cell lines. *J Gene Med* 2007, 9(2):122–135.
- [52] Kariko K, Weissman D: Naturally occurring nucleoside modifications suppress the immunostimulatory activity of RNA: implication for therapeutic RNA development. *Curr Opin Drug Discov Dev* 2007, 10(5):523–532.
- [53] Kariko K, Muramatsu H, Welsh FA, Ludwig J, Kato H, Akira S, Weissman D: Incorporation of pseudouridine into mRNA yields superior nonimmunogenic vector with increased translational capacity and biological stability. *Mol Ther* 2008, 16(11):1833–1840.
- [54] Anderson BR, Muramatsu H, Nallagatla SR, Bevilacqua PC, Sansing LH, Weissman D, Kariko K: Incorporation of pseudouridine into mRNA enhances translation by diminishing PKR activation. *Nucleic Acids Res* 2010, 38(17):5884–5892.
- [55] Hornung V, Ellegast J, Kim S, Brzozka K, Jung A, Kato H, Poeck H, Akira S, Conzelmann KK, Schlee M, et al.: 5'-Triphosphate RNA is the ligand for RIG-I. *Science* 2006, 314(5801):994–997.
- [56] Kariko K, Muramatsu H, Ludwig J, Weissman D: Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, protein-encoding mRNA. *Nucleic Acids Res* 2011, 39(21):e142.
- [57] Dyer KD, Rosenberg HF: The RNase a superfamily: generation of diversity and innate host defense. *Mol Diversity* 2006, 10(4):585–597.
- [58] McKenzie R, Fried MW, Sallie R, Conjeevaram H, Di Bisceglie AM, Park Y, Savarese B, Kleiner D, Tsokos M, Luciano C, et al.: Hepatic failure and lactic acidosis due to

- fialuridine (FIAU), an investigational nucleoside analogue for chronic hepatitis B. *N Engl J Med* 1995, 333(17):1099–1105.
- [59] Lewis W: Defective mitochondrial DNA replication and NRTIs: pathophysiological implications in AIDS cardiomyopathy. *Am J Physiol Heart Circ Physiol* 2003, 284(1):H1–H9.
- [60] Yin H, Kanasty RL, Eltoukhy AA, Vegas AJ, Dorkin JR, Anderson DG: Non-viral vectors for gene-based therapy. *Nat Rev Genet* 2014, 15(8):541–555.
- [61] Agapov EV, Frolov I, Lindenbach BD, Pragai BM, Schlesinger S, Rice CM: Noncytopathic Sindbis virus RNA vectors for heterologous gene expression. *Proc Natl Acad Sci U S A* 1998, 95(22):12989–12994.
- [62] Bitzer M, Armeanu S, Lauer UM, Neubert WJ: Sendai virus vectors as an emerging negative-strand RNA viral vector system. *J Gene Med* 2003, 5(7):543–553.
- [63] Baum C, Kustikova O, Modlich U, Li Z, Fehse B: Mutagenesis and oncogenesis by chromosomal insertion of gene transfer vectors. *Hum Gene Ther* 2006, 17(3):253–263.
- [64] Bessis N, GarciaCozar FJ, Boissier MC: Immune responses to gene therapy vectors: influence on vector function and effector mechanisms. *Gene Ther* 2004, 11 Suppl 1:S10–S17.
- [65] Bouard D, Alazard-Dany D, Cosset FL: Viral vectors: from virology to transgene expression. *Br J Pharmacol* 2009, 157(2):153–165.
- [66] Thomas CE, Ehrhardt A, Kay MA: Progress and problems with the use of viral vectors for gene therapy. *Nat Rev Genet* 2003, 4(5):346–358.
- [67] Pack DW, Hoffman AS, Pun S, Stayton PS: Design and development of polymers for gene delivery. *Nat Rev Drug Discov* 2005, 4(7):581–593.
- [68] Mintzer MA, Simanek EE: Nonviral vectors for gene delivery. *Chem Rev* 2009, 109(2):259–302.
- [69] Van Tendeloo VF, Ponsaerts P, Lardon F, Nijs G, Lenjou M, Van Broeckhoven C, Van Bockstaele DR, Berneman ZN: Highly efficient gene delivery by mRNA electroporation in human hematopoietic cells: superiority to lipofection and passive pulsing of mRNA and to electroporation of plasmid cDNA for tumor antigen loading of dendritic cells. *Blood* 2001, 98(1):49–56.
- [70] Sohn RL, Murray MT, Schwarz K, Nyitray J, Purray P, Franko AP, Palmer KC, Diebel LN, Dulchavsky SA: *In-vivo* particle mediated delivery of mRNA to mammalian tissues: ballistic and biologic effects. *Wound Repair Regen* 2001, 9(4):287–296.
- [71] Zhdanov RI, Podobed OV, Vlassov VV: Cationic lipid-DNA complexes-lipoplexes-for gene transfer and therapy. *Bioelectrochemistry* 2002, 58(1):53–64.

- [72] Zou S, Scarfo K, Nantz MH, Hecker JG: Lipid-mediated delivery of RNA is more efficient than delivery of DNA in non-dividing cells. *Int J Pharm* 2010, 389(1–2):232–243.
- [73] Luo D, Saltzman WM: Synthetic DNA delivery systems. *Nat Biotechnol* 2000, 18(1):33–37.
- [74] Tros de Ilarduya C, Duzgunes N: Efficient gene transfer by transferrin lipoplexes in the presence of serum. *Biochim Biophys Acta* 2000, 1463(2):333–342.
- [75] Noble GT, Stefanick JF, Ashley JD, Kiziltepe T, Bilgicer B: Ligand-targeted liposome design: challenges and fundamental considerations. *Trends Biotechnol* 2014, 32(1):32–45.
- [76] Allen TM, Cullis PR: Liposomal drug delivery systems: from concept to clinical applications. *Adv Drug Deliv Rev* 2013, 65(1):36–48.
- [77] Andresen TL, Jensen SS, Jorgensen K: Advanced strategies in liposomal cancer therapy: problems and prospects of active and tumor specific drug release. *Prog Lipid Res* 2005, 44(1):68–97.
- [78] Ruoslahti E: Peptides as targeting elements and tissue penetration devices for nanoparticles. *Adv Mater* 2012, 24(28):3747–3756.
- [79] Matsui A, Uchida S, Ishii T, Itaka K, Kataoka K: Messenger RNA-based therapeutics for the treatment of apoptosis-associated diseases. *Sci Rep* 2015, 5:15810.
- [80] Kariko K, Muramatsu H, Keller JM, Weissman D: Increased erythropoiesis in mice injected with submicrogram quantities of pseudouridine-containing mRNA encoding erythropoietin. *Mol Ther* 2012, 20(5):948–953.
- [81] Mays LE, Ammon-Treiber S, Mothes B, Alkhaled M, Rottenberger J, Muller-Hermelink ES, Grimm M, Mezger M, Beer-Hammer S, von Stebut E, et al.: Modified Foxp3 mRNA protects against asthma through an IL-10-dependent mechanism. *J Clin Invest* 2013, 123(3):1216–1228.
- [82] Granstein RD, Ding W, Ozawa H: Induction of anti-tumor immunity with epidermal cells pulsed with tumor-derived RNA or intradermal administration of RNA. *J Invest Dermatol* 2000, 114(4):632–636.
- [83] Kreiter S, Selmi A, Diken M, Koslowski M, Britten CM, Huber C, Tureci O, Sahin U: Intranodal vaccination with naked antigen-encoding RNA elicits potent prophylactic and therapeutic antitumoral immunity. *Cancer Res* 2010, 70(22):9031–9040.
- [84] Azarmi S, Roa WH, Lobenberg R: Targeted delivery of nanoparticles for the treatment of lung diseases. *Adv Drug Deliv Rev* 2008, 60(8):863–875.
- [85] Johler SM, Rejman J, Guan S, Rosenecker J: Nebulisation of IVT mRNA complexes for intrapulmonary administration. *PLoS One* 2015, 10(9):e0137504.
- [86] Sheridan C: Gene therapy finds its niche. *Nat Biotechnol* 2011, 29(2):121–128.

- [87] Das SK, Menezes ME, Bhatia S, Wang XY, Emdad L, Sarkar D, Fisher PB: Gene therapies for cancer: strategies, challenges and successes. *J Cell Physiol* 2015, 230(2):259–271.
- [88] Stoller JK, Fallat R, Schluchter MD, O'Brien RG, Connor JT, Gross N, O'Neil K, Sandhaus R, Crystal RG: Augmentation therapy with α 1-antitrypsin: patterns of use and adverse events. *Chest* 2009, 136(5 Suppl):e30.
- [89] Silverman EK, Sandhaus RA: Clinical practice. α 1-Antitrypsin deficiency. *N Engl J Med* 2009, 360(26):2749–2757.
- [90] Chill L, Trinh L, Azadi P, Ishihara M, Sonon R, Karnaukhova E, Ophir Y, Golding B, Shiloach J: Production, purification, and characterization of human α 1 proteinase inhibitor from *Aspergillus niger*. *Biotechnol Bioeng* 2009, 102(3):828–844.
- [91] Courtney M, Buchwalder A, Tessier LH, Jaye M, Benavente A, Balland A, Kohli V, Lathe R, Tolstoshev P, Lecocq JP: High-level production of biologically active human α 1-antitrypsin in *Escherichia coli*. *Proc Natl Acad Sci U S A* 1984, 81(3):669–673.
- [92] Kwon KS, Song M, Yu MH: Purification and characterization of α 1-antitrypsin secreted by recombinant yeast *Saccharomyces diastaticus*. *J Biotechnol* 1995, 42(3):191–195.
- [93] Huang J, Sutliff TD, Wu L, Nandi S, Bengel K, Terashima M, Ralston AH, Drohan W, Huang N, Rodriguez RL: Expression and purification of functional human α 1-antitrypsin from cultured plant cells. *Biotechnol Prog* 2001, 17(1):126–133.
- [94] Ross D, Brown T, Harper R, Pamarthi M, Nixon J, Bromirski J, Li CM, Ghali R, Xie H, Medvedeff G, et al.: Production and characterization of a novel human recombinant α 1-antitrypsin in PER.C6 cells. *J Biotechnol* 2012, 162(2–3):262–273.
- [95] Niklas J, Priesnitz C, Rose T, Sandig V, Heinzle E: Metabolism and metabolic burden by α 1-antitrypsin production in human AGE1.HN cells. *Metab Eng* 2013, 16:103–114.
- [96] Blanchard V, Liu X, Eigel S, Kaup M, Rieck S, Janciauskiene S, Sandig V, Marx U, Walden P, Tauber R, et al.: N-glycosylation and biological activity of recombinant human α 1-antitrypsin expressed in a novel human neuronal cell line. *Biotechnol Bioeng* 2011, 108(9):2118–2128.
- [97] Jirikowski GF, Sanna PP, Maciejewski-Lenoir D, Bloom FE: Reversal of diabetes insipidus in Brattleboro rats: intrahypothalamic injection of vasopressin mRNA. *Science* 1992, 255(5047):996–998.
- [98] Mahiny AJ, Dewerth A, Mays LE, Alkhaled M, Mothes B, Malaeksefat E, Loretz B, Rottenberger J, Brosch DM, Reautschnig P, et al.: *In vivo* genome editing using nuclease-encoding mRNA corrects SP-B deficiency. *Nat Biotechnol* 2015, 33(6):584–586.
- [99] Michel T, Kankura A, Salinas Medina ML, Kurz J, Behring A, Avci-Adali M, Nolte A, Schlensak C, Wendel HP, Krajewski S: *In vitro* evaluation of a novel mRNA-based therapeutic strategy for the treatment of patients suffering from α 1-antitrypsin deficiency. *Nucleic Acids Ther* 2015, 25(5):235–244.